

Psychological investigation of the structure of paranoia in a non-clinical population

DANIEL FREEMAN, PHILIPPA A. GARETY, PAUL E. BEBBINGTON, BENJAMIN SMITH, REBECCA ROLLINSON, DAVID FOWLER, ELIZABETH KUIPERS, KATARZYNA RAY and GRAHAM DUNN

Background Previous studies of paranoia have assessed only limited numbers of paranoid thoughts, and have not considered the experience from a multidimensional perspective or examined the relationship between different suspicious thoughts.

Aims To assess a wide range of paranoid thoughts multidimensionally and examine their distribution, to identify the associated coping strategies and to examine social–cognitive processes and paranoia.

Method Six questionnaire assessments were completed by 1202 individuals using the internet.

Results Paranoid thoughts occurred regularly in approximately a third of the group. Increasing endorsement of paranoid thoughts was characterised by the recruitment of rarer and odder ideas. Higher levels of paranoia were associated with emotional and avoidant coping, less use of rational and detached coping, negative attitudes to emotional expression, submissive behaviours and lower social rank.

Conclusions Suspiciousness is common and there may be a hierarchical arrangement of such thoughts that builds on common emotional concerns.

Declaration of interest None.

It is possible that paranoid ideation is almost as common as symptoms of anxiety and depression (e.g. van Os & Verdoux, 2003; Johns *et al*, 2004). For many people, thoughts that friends, acquaintances or strangers might be hostile, or deliberately watching them, appear to be an everyday occurrence. This may parallel the findings of earlier studies that a level of obsessive thinking is normal in the general population (Rachman & de Silva, 1978). In one way this is not surprising, because being wary of the intentions of others is adaptive in some situations. At a cultural level, a fear of others is variably incorporated into the political and social climate. In general such thoughts are not a clinical problem, becoming so only when they are excessive, exaggerated or unfounded, and cause distress. Paranoid thoughts in non-clinical populations are phenomena of interest in their own right and may inform our understanding of delusions.

This study provides information on the frequency in a non-clinical sample of paranoid ideation, and the associated levels of conviction and distress. Such information can be useful to present to patients in the clinical setting, but no comparable research examining a wide range of paranoid thoughts, or considering such thoughts from a multidimensional perspective, has hitherto been published. There is no published evidence on, for example, the weekly frequency of paranoia in the general population. We predicted that the distribution of suspicious thoughts would be similar to that of affective symptoms, with many people having a few suspicious thoughts and a few people having many (Melzer *et al*, 2002). Moreover, as with affective symptoms, increasing symptom counts will be characterised by the recruitment of rarer and odder ideas (Sturt, 1981). There may be a hierarchy of paranoid thoughts. The study also had the aim of identifying how individuals in the general population cope

with paranoid thoughts. We wished to identify the coping strategies that were associated with the most and the least distress. Finally, we examined potential connections between paranoia and three social–cognitive processes: attitudes to emotional expression, social comparison and submissive behaviours.

METHOD

Participants and procedure

An anonymous internet survey was considered to provide a safe environment for survey participants to disclose suspicious thoughts. Internet research has been found to reach the same conclusions as laboratory-based studies (Birnbaum, 2001). Students at King's College London, the University of East Anglia and University College London were e-mailed the address of a website where they could take part in a survey of 'everyday worries about others'. The study had received the approval of local research ethics committees. Completion of each questionnaire was timed, and one submission completed too quickly (<45 s) was deleted. Paranoia Checklist questionnaires with more than five pieces of missing data (i.e. completion rate less than 90%) were not included; consequently, the percentage of the Paranoia Checklist data that were prorated was minimal (<0.5%). The final sample comprised 1202 people. After providing demographic information, participants were presented with six questionnaires.

Questionnaires

Paranoia Scale

The 20-item, self-report Paranoia Scale (Fenigstein & Vanable, 1992) was developed to measure paranoia in college students. Each item is rated on a five-point scale (1 not at all applicable, 5 extremely applicable). Scores can range from 20 to 100, with the higher scores indicating greater paranoid ideation. It is the most widely used dimensional measure of paranoia. However, the scale contains many items that are not clearly persecutory (e.g. 'My parents and family find more fault with me than they should') and does not provide an estimate of the frequency or distress of paranoid thoughts. The Paranoia Checklist was therefore developed specifically for this study.

Paranoia Checklist

The Paranoia Checklist was devised to investigate paranoid thoughts of a more clinical nature than those assessed in the Paranoia Scale and to provide a multi-dimensional assessment of paranoid ideation. The checklist has 18 items, each rated on a five-point scale for frequency, degree of conviction, and distress. We report the convergent validity of the Paranoia Checklist in relation to the Paranoia Scale in the results section.

Coping Styles Questionnaire

The Coping Styles Questionnaire (CSQ; Roger *et al*, 1993) builds upon the Ways of Coping Checklist (Folkman & Lazarus, 1980), and was validated with a UK student sample. The questionnaire comprises 60 coping strategies rated on a four-point frequency scale. Participants were asked to complete the questionnaire for how they typically react to the worries assessed in the Paranoia Checklist. There are four factors: rational coping, detached coping, emotional coping and avoidance coping. The former two factors are considered by the questionnaire developers as adaptive and the latter two as maladaptive.

Attitudes to Emotional Expression Questionnaire

On the Attitudes to Emotional Expression Questionnaire (Joseph *et al*, 1994) respondents were asked to rate how much they agree on a five-point scale (1 agree very much, 2 agree slightly, 3 neutral, 4 disagree slightly, 5 disagree very much) with four attitudes to emotional expression (e.g. 'I think you should always keep your feelings under control'). Higher scores indicate more positive attitudes to emotional expression.

Social Comparison Scale

On the Social Comparison Scale (Gilbert & Allan, 1994) participants rate, by selecting a number between 1 and 10, whether they generally feel in relation to others: inferior–superior; less competent–more competent; less likeable–likeable; more reserved–less reserved; left out–accepted. Higher scores indicate higher perceived social rank.

Submissive Behaviours Scale

The Submissive Behaviours Scale (Allan & Gilbert, 1997) is a 16-item scale assessing a number of behaviours considered as submissiveness (e.g. 'I agree that I am wrong, even though I know I'm not'). Each behaviour is rated on a five-point scale (0 never, 4 always). Higher scores indicate greater use of submissive behaviours.

Analysis

Analyses were conducted using the Statistical Package for the Social Sciences, SPSS for Windows, version 11.0 (SPSS, 2001). Significance test results are quoted as two-tailed probabilities.

RESULTS

Demographic data

There were more women ($n=821$) than men ($n=371$) among the respondents. Although the World Wide Web was once considered predominantly a male preserve, studies have found that more women than men participate in online psychological studies (Birnbaum, 2001). The average age was 23.0 years (s.d.=6.1, range 17–61, interquartile range 5). The respondents reported their ethnicity predominantly as White ($n=1001$), followed by Asian ($n=98$) and 'other' ($n=70$). There were few Black African ($n=9$) and African–Caribbean ($n=9$) respondents.

Reliability and validity of the Paranoia Checklist

Cronbach's α for each of the three dimensions of the Paranoia Checklist was 0.9 or above, indicating excellent internal reliability. As would be expected, time taken to complete the Paranoia Checklist had a small positive correlation with a total score for frequency, conviction and distress ($r=0.10$, $P<0.001$). The Paranoia Scale was completed by 1016 of the Paranoia Checklist respondents. The mean Paranoia Scale score of the total group was 42.7 (s.d.=14.3), which is comparable with that reported by Fenigstein & Venable (1992). There was convergent validity of the checklist with the Paranoia Scale: higher Paranoia Scale scores correlated with Paranoia Checklist frequency ($r=0.71$, $P<0.001$), conviction ($r=0.62$, $P<0.001$) and distress scores ($r=0.58$, $P<0.001$).

Prevalence of thoughts with a paranoid content

The frequencies, conviction and distress associated with the suspicious thoughts assessed in the Paranoia Checklist are displayed in Tables 1–3. There was appreciable endorsement of the checklist items. The 1-week prevalence of the individual thoughts ranged from 3% ('I can detect messages about me in the press/TV/radio') to 52% ('I need to be on my guard against others') (Table 1). The mean frequency score was 11.9 (s.d.=10.5, range 0–64; 25th percentile 4.0, 50th percentile 9.0, 75th percentile 16.0). Between 2% and 7% of participants adhered to individual thoughts with a level of absolute conviction (Table 2). If we consider levels of belief of 'somewhat' or greater, there is more variation between the individual items (4–56%). The mean conviction score was 16.7 (s.d.=12.1, range 0–72; 25th percentile 8.0, 50th percentile 14.0, 75th percentile 22.0). Between 1% and 7% of participants found individual thoughts very distressing (Table 3). Again, there was more variation between the thoughts if distress was taken as 'at least somewhat distressing' or greater (3–42%). The mean distress score was 14.6 (s.d.=12.2, range 0–70; 25th percentile 5.0, 50th percentile 12.0, 75th percentile 21.0).

The different dimensions of the Paranoia Checklist were positively correlated. Frequency scores were correlated with conviction ($r=0.75$, $P<0.001$) and distress ($r=0.66$, $P<0.001$), and conviction and distress scores were also positively correlated ($r=0.65$, $P<0.001$). There were no differences between men and women in the frequency ($t=0.66$, d.f.=1190, $P=0.51$) or conviction ($t=1.03$, d.f.=1190, $P=0.30$) with which paranoid thoughts were experienced. Females did report a significantly higher level of distress associated with the thoughts ($t=-2.72$, d.f.=1190, $P=0.007$, mean difference = -2.07, 95% CI -3.60 to -0.58), although it can be seen that this difference is very small.

In Tables 4 and 5 the levels of conviction and distress associated with each suspicious thought are reported for the individuals experiencing such ideas at least weekly. Here it can be seen that the rarer and more implausible paranoid items (e.g. 'There is a possibility of a conspiracy against me') are held with the strongest levels of conviction and associated with

Table 1 Frequency ('How often have you had the thought?') (n=1202)

	Rarely (%)	Once a month (%)	Once a week (%)	Several times a week (%)	At least once a day (%)	Weekly (%)
I need to be on my guard against others	31	17	21	21	10	52
There might be negative comments being circulated about me	35	24	21	14	7	42
People deliberately try to irritate me	57	17	15	8	4	27
I might be being observed or followed	67	14	8	7	4	19
People are trying to make me upset	72	16	7	4	1	12
People communicate about me in subtle ways	52	22	14	9	3	26
Strangers and friends look at me critically	29	23	21	18	9	48
People might be hostile towards me	45	27	16	9	4	29
Bad things are being said about me behind my back	45	25	15	11	4	30
Someone I know has bad intentions towards me	71	16	6	4	2	12
I have a suspicion that someone has it in for me	83	9	4	2	2	8
People would harm me if given an opportunity	83	9	4	2	2	8
Someone I don't know has bad intentions towards me	82	10	3	3	2	8
There is a possibility of a conspiracy against me	90	5	2	1	2	5
People are laughing at me	41	26	19	9	6	34
I am under threat from others	76	13	5	3	2	10
I can detect coded messages about me in the press/TV/radio	96	2	1	1	1	3
My actions and thoughts might be controlled by others	81	10	3	3	2	8

Table 2 Conviction ('How strongly do you believe it?') (n=1202)

	Do not believe it (%)	Believe it a little (%)	Believe it somewhat (%)	Believe it a lot (%)	Absolutely believe it (%)	Somewhat or greater (%)
I need to be on my guard against others	10	34	34	17	5	56
There might be negative comments being circulated about me	13	41	28	13	5	46
People deliberately try to irritate me	36	33	17	10	5	32
I might be being observed or followed	51	29	11	7	2	20
People are trying to make me upset	51	29	11	5	3	19
People communicate about me in subtle ways	37	33	19	8	3	30
Strangers and friends look at me critically	19	31	27	16	7	50
People might be hostile towards me	25	37	24	9	6	39
Bad things are being said about me behind my back	26	39	20	11	5	36
Someone I know has bad intentions towards me	51	29	10	5	5	20
I have a suspicion that someone has it in for me	67	18	8	3	4	15
People would harm me if given an opportunity	67	19	8	3	3	14
Someone I don't know has bad intentions towards me	72	16	6	3	3	12
There is a possibility of a conspiracy against me	83	10	3	2	3	8
People are laughing at me	30	35	19	10	5	34
I am under threat from others	65	20	9	3	3	15
I can detect coded messages about me in the press/TV/radio	93	3	1	1	2	4
My actions and thoughts might be controlled by others	77	12	5	2	4	11

the most distress. This is confirmed by high negative correlations between the frequency of (at least weekly) endorsement of questionnaire items and the percentage of people who believed the thought absolutely

($n=18$; $r=-0.74$, $P<0.001$) and with the percentage of people who found the thought very distressing ($n=18$; $r=-0.75$, $P<0.001$). In other words, the less frequently experienced thoughts were held

with proportionately more conviction and distress.

To examine whether people who endorsed the rarer items also endorsed the more common suspicious thoughts with

Table 3 Distress ('How upsetting is it for you?') (n=1202)

	Not distressing (%)	A little distressing (%)	Somewhat distressing (%)	Moderately distressing (%)	Very distressing (%)	At least somewhat distressing (%)
I need to be on my guard against others	33	36	18	11	3	32
There might be negative comments being circulated about me	25	34	20	16	6	42
People deliberately try to irritate me	48	30	11	7	4	22
I might be being observed or followed	55	23	11	7	3	21
People are trying to make me upset	53	23	13	7	5	25
People communicate about me in subtle ways	51	29	12	5	2	19
Strangers and friends look at me critically	34	31	18	12	5	35
People might be hostile towards me	35	34	18	10	4	32
Bad things are being said about me behind my back	34	31	17	11	7	35
Someone I know has bad intentions towards me	54	23	12	7	5	24
I have a suspicion that someone has it in for me	66	18	8	4	3	15
People would harm me if given an opportunity	67	15	7	4	6	17
Someone I don't know has bad intentions towards me	72	16	6	3	3	12
There is a possibility of a conspiracy against me	82	9	4	3	3	10
People are laughing at me	43	28	15	9	5	29
I am under threat from others	66	18	8	5	3	16
I can detect coded messages about me in the press/TV/radio	96	2	1	1	1	3
My actions and thoughts might be controlled by others	82	10	4	2	2	8

Table 4 Level of conviction for those who experienced the thought at least weekly

	n	Do not believe it (%)	Believe it a little (%)	Believe it somewhat (%)	Believe it a lot (%)	Absolutely believe it (%)
I need to be on my guard against others	621	1	19	44	27	9
There might be negative comments being circulated about me	499	1	23	43	23	9
People deliberately try to irritate me	306	3	24	37	23	12
I might be being observed or followed	230	4	27	36	25	8
People are trying to make me upset	148	1	25	38	25	11
People communicate about me in subtle ways	315	1	21	44	25	10
Strangers and friends look at me critically	574	0	16	42	30	13
People might be hostile towards me	343	0	18	47	21	14
Bad things are being said about me behind my back	360	3	21	39	25	13
Someone I know has bad intentions towards me	155	2	21	33	21	23
I have a suspicion that someone has it in for me	97	3	21	28	21	28
People would harm me if given an opportunity	94	0	5	47	25	23
Someone I don't know has bad intentions towards me	91	0	20	37	18	25
There is a possibility of a conspiracy against me	58	5	22	22	24	26
People are laughing at me	393	1	22	38	25	13
I am under threat from others	131	2	16	45	22	15
I can detect coded messages about me in the press/TV/radio	29	0	17	28	21	35
My actions and thoughts might be controlled by others	100	2	18	36	20	24

higher conviction and distress, we split the sample into those who endorsed at least one rare item (items with frequency less than 10%) (n=277) and those who did

not endorse a rare item (n=925). The two groups were then compared on levels of conviction and distress for the eight most common items (endorsement rates of over

20%). Each one of these comparisons was made only for the individuals in the two groups who had endorsed the item (i.e. experienced the thought at least weekly).

Table 5 Level of distress for those who experienced the thought at least weekly

	<i>n</i>	Not distressing (%)	A little distressing (%)	Somewhat distressing (%)	Moderately distressing (%)	Very distressing (%)
I need to be on my guard against others	621	19	34	26	16	5
There might be negative comments being circulated about me	499	10	28	25	26	12
People deliberately try to irritate me	306	18	34	25	15	9
I might be being observed or followed	230	16	29	25	18	11
People are trying to make me upset	148	5	22	35	17	22
People communicate about me in subtle ways	315	19	32	27	16	6
Strangers and friends look at me critically	574	13	29	27	21	10
People might be hostile towards me	343	9	29	30	21	11
Bad things are being said about me behind my back	360	8	22	29	23	19
Someone I know has bad intentions towards me	155	9	23	30	16	21
I have a suspicion that someone has it in for me	97	11	23	28	13	25
People would harm me if given an opportunity	94	6	14	33	15	32
Someone I don't know has bad intentions towards me	91	12	22	26	20	20
There is a possibility of a conspiracy against me	58	9	24	22	21	24
People are laughing at me	393	17	25	25	19	14
I am under threat from others	131	8	19	30	28	15
I can detect coded messages about me in the press/TV/radio	29	24	17	21	17	21
My actions and thoughts might be controlled by others	100	12	28	24	17	19

The rarer item group had significantly higher conviction rates for seven of the eight common suspicious thoughts and significantly higher distress levels for five of the eight common suspicious thoughts ($P < 0.05$). It seems that individuals with the rarer thoughts were also experiencing the commoner thoughts more strongly.

We predicted that suspiciousness in the general population would have a profile similar to that of affective symptoms. First, there would be a single population distribution rather than evidence of a bimodal distribution (i.e. between 'clinical paranoia' and 'non-clinical paranoia'). Second, the rarer ideas would be associated with the presence of many other suspicions; put another way, the relationship between rare symptoms and common symptoms would be non-reflexive, in that the former would be more predictive of the latter than *vice versa*. The total number of checklist items endorsed by each person was first calculated (endorsement referring to weekly occurrence or above). The count of suspicious thoughts could therefore range from 0 to 18. The distribution of the count is displayed in Fig. 1. It can be seen that the suspicious thought count follows a single continuous model (Melzer *et al*, 2002). The distribution closely fits an exponential

curve. To examine whether the rarer thoughts were associated with a higher rate of endorsement of other checklist items, the mean difference for the suspicious thought count was calculated between those with and those without each suspicious thought (correcting for the contribution due to that

item; Sturt, 1981). The mean difference (i.e. the excess of endorsement associated with each item) was significantly associated with the frequency of item endorsement ($n=18$; $r=-0.75$, $P < 0.001$). Thus, the rarer checklist items were associated with a higher total score than were the




Fig. 1 Percentage of the study population v. total number of Paranoia Checklist items endorsed, with fitted exponential curve.

Table 6 Coping strategies and paranoia (n=1046)

Type of coping	Paranoia Checklist					
	Frequency		Conviction		Distress	
	r	P	r	P	r	P
Emotional	0.56	<0.001	0.42	<0.001	0.55	<0.001
Avoidant	0.33	<0.001	0.24	<0.001	0.28	<0.001
Detached	-0.25	<0.001	-0.14	<0.001	-0.33	<0.001
Rational	-0.17	<0.001	-0.06	0.06	-0.20	<0.001

more common ones. For example, endorsing the item ‘There might be negative comments being circulated about me’ (frequency 42%) was associated with endorsing 3.9 other checklist items in comparison with not endorsing the item. Endorsing the item ‘There is a possibility of a conspiracy against me’ (frequency 5%) was associated with endorsing 7.0 other checklist items in comparison with not endorsing the item.

Coping with paranoid thoughts

A total of 1046 participants also completed the CSQ. Higher levels of emotional and avoidant coping were associated with higher levels of paranoia (Table 6). In contrast, higher levels of detached coping were associated with lower levels of paranoia. Higher levels of rational coping were associated with lower levels of paranoia frequency and distress, but not significantly with

paranoia conviction. In Table 7 we highlight the coping strategies most strongly correlated with paranoia frequency.

Social-cognitive processes and paranoid thoughts

There were significant but generally modest associations between the social-cognitive processes and the dimensions of paranoia. Negative attitudes to emotional expression, lower social comparison (particularly feeling left out) and greater use of submissive behaviours were significantly associated with greater paranoia (Table 8).

DISCUSSION

Study limitations

There are methodological constraints that need to be acknowledged when interpreting our results. Foremost, the sample was not epidemiologically representative. It was self-selected, restricted to university students and recruited by e-mail. The gender ratio was certainly skewed, perhaps

Table 7 Coping strategies most associated with high and low frequencies of paranoia (n=1046)

Coping strategy	Paranoia Checklist		
	Frequency (r)	Conviction (r)	Distress (r)
Presence associated with more frequent paranoia			
Become lonely or isolated	0.486	0.329	0.409
Feel that no one understands	0.471	0.354	0.397
Feel worthless and unimportant	0.456	0.330	0.444
Become miserable or distressed	0.445	0.315	0.430
Feel helpless – there’s nothing you can do about it	0.415	0.270	0.399
Criticise or blame myself	0.403	0.327	0.402
Avoid family or friends in general	0.390	0.292	0.312
Feel overpowered and at the mercy of the situation	0.381	0.304	0.417
Stop doing hobbies or interests	0.340	0.250	0.348
Daydream about times in the past when things were better	0.337	0.260	0.313
Presence associated with less frequent paranoia			
Do not see the problem or situation as a threat	-0.327	-0.220	-0.325
See the situation for what it is and nothing more	-0.266	-0.147	-0.321
Try to find the positive side to the situation	-0.248	-0.176	-0.275
Have presence of mind when dealing with the problem or situation	-0.246	-0.155	-0.281
Feel completely clear-headed about the whole thing	-0.237	-0.163	-0.291
Be realistic in my approach to the situation	-0.229	-0.136	-0.220
See the problem as something separate from myself so I can deal with it	-0.199	-0.122	-0.218
Keep reminding myself about the good things about myself	-0.198	-0.152	-0.190
Get things into proportion – nothing is really that important	-0.194	-0.135	-0.269
Just take nothing personally	-0.181	-0.148	-0.289

All significant at P < 0.001.

Table 8 Social cognitive processes and paranoia

	n	Paranoia Checklist					
		Frequency		Conviction		Distress	
		r	P	r	P	r	P
Attitudes to emotional expression	1049	0.291	<0.001	0.235	<0.001	0.161	<0.001
Social comparison							
Mean score	1035	-0.265	<0.001	-0.193	<0.001	-0.270	<0.001
Inferior–superior	966	-0.149	<0.001	-0.091	0.005	-0.179	<0.001
Less competent–more competent	980	-0.117	<0.001	-0.089	0.005	-0.190	<0.001
Less likeable–likeable	934	-0.038	0.251	-0.051	0.001	-0.010	0.763
More reserved–less reserved	938	-0.092	0.005	-0.058	0.074	-0.081	0.013
Left out–accepted	970	-0.374	<0.001	-0.271	<0.001	-0.333	<0.001
Submissive behaviour	1027	0.392	<0.001	0.295	<0.001	0.385	<0.001

in consequence of our methods. People who self-select for questionnaires of this type may be more prone to psychological disturbance, or the stigma of appearing so might skew the sample in the opposite direction. Thus, our investigation in a selected group indicates a need for more elaborate and more truly epidemiological studies. There are also issues concerning whether the experiences assessed are actually unfounded; questionnaire studies may include an unknown proportion of paranoia that is realistic and therefore well judged and appropriate. It is also unknown whether any of the participants had received treatment for a psychiatric disorder, and what the level of substance use was in the group. The study was also limited by the cross-sectional questionnaire design, which limits the conclusions that can be drawn concerning the causal direction of the relationships between associated variables.

Hierarchy of paranoia


Our survey clearly indicates that suspicious thoughts are a weekly occurrence for many people: 30–40% of the respondents had ideas that negative comments were being circulated about them and 10–30% had persecutory thoughts, with thoughts of mild threat (e.g. 'People deliberately try to irritate me') being more common than severe threat (e.g. 'Someone has it in for me'). In contrast, only a small proportion (approximately 5%) of respondents endorsed the checklist items that were the most improbable (e.g. that there was a conspiracy). Nevertheless, the rarer and odder suspicions – characteristic of clinical

presentations – occurred in tandem with the more common and plausible experiences. The rarer the thought, then the higher the total score indicated by its presence. There has been no previous examination of paranoia in this way.

The findings indicate a hierarchy of paranoia (Fig. 2): the most common type of suspiciousness is that of a social anxiety or interpersonal worry theme; ideas of reference build upon these sensitivities; persecutory thoughts are closely associated with the attributions of significance; as the severity of the threatened harm increases, the less common the thought; and suspiciousness involving severe harm and organisations and conspiracy is at the top of the hierarchy. The implication is that

severe paranoia may build upon common emotional concerns, consistent with a recent cognitive model of persecutory delusions (Freeman *et al.*, 2002; Freeman & Garety, 2004). The interesting questions therefore concern the identification of the additional factors that contribute to the development of severe paranoia and whether there are qualitative shifts in experience at the top end of the hierarchy (note that individuals at the higher end of the hierarchy tended to endorse all their suspicious thoughts with high levels of conviction and distress). The survey findings also indicate that there is a continuous (exponential) distribution of total number of suspicious thoughts in the general population, although the thoughts appear in a hierarchical arrangement. No distinct sub-population was identified. This therefore demonstrates correspondence to common mental health disorders such as depression and anxiety.

Interestingly, the ideation captured in this survey did not seem to be restricted to passing thoughts that were dismissed almost in the same instant that they occurred. Approximately 10–20% of the survey respondents held paranoid ideation with strong conviction and significant distress. It is likely that the survey identified a significant group of people who were having distressing experiences that they managed on their own. We believe that there is a reticence in the general population about discussing the occurrence of suspicious thoughts, partly arising from the negative connotations associated with the term

**Fig. 2** The paranoia hierarchy.

paranoia and a lack of recognition of how common these experiences actually are. The provision of the type of information obtained in this survey may help to normalise paranoia and set the stage for making the experience understandable. This is an important element in the development of alternative explanations of experiences (Freeman *et al*, 2004).

Coping with paranoid thoughts

If paranoia is an everyday phenomenon, which many people manage well, then it provides an opportunity to gain clinically useful information on optimal ways of coping. More frequent and distressing paranoia was associated with becoming isolated, giving up activities, and feelings of powerlessness and depression. Conversely, less frequent paranoia was associated with not catastrophising and by gaining sufficient (metacognitive) distance to consider the situation dispassionately. More broadly, coping with paranoia may resemble coping with other stressful or negative events: rational (or task-oriented) coping and detachment from the situation are more helpful than emotional or avoidant coping. It is not clear to what extent poor coping encourages paranoia, and to what extent strong paranoia interferes with effective coping.

Social-cognitive factors

There has been a re-emergence of the study of the influence of social factors on psychosis by examining their impact at the cognitive level of explanation (Garety *et al*, 2001). In our survey associations were found between paranoia and social-cognitive processes that could plausibly exacerbate suspiciousness. Thus, we found evidence that not expressing feelings to others may increase suspiciousness. This follows early research on the psychological consequences of the *Herald of Free Enterprise* ferry disaster, which indicated that negative attitudes to emotional expression in survivors were associated with higher levels of anxiety (Joseph *et al*, 1997).

There was a significant association of paranoia with submissive behaviours. As Allan & Gilbert (1997) note, people who have difficulties in asserting themselves – which these authors conceptualise within an evolutionary framework as having low dominance and inferior social rank – can be vulnerable to a number of psychological problems. Further, these authors report that submissiveness is associated with

CLINICAL IMPLICATIONS

- Having suspicious thoughts is a common experience and provision of this information may help reduce patient distress.
- Feelings of hopelessness and lack of control may contribute to the occurrence of more suspicious thoughts, whereas gaining distance from such thoughts and evaluating them may reduce such experiences.
- Not talking to others about suspicious thoughts, feeling vulnerable and behaving timidly with others may be factors in the development of paranoia.

LIMITATIONS

- An epidemiologically representative sample was not recruited.
- The group mainly comprised young adults who may have higher rates of suspiciousness.
- Only cross-sectional associations between paranoia, coping strategies and social-cognitive processes were examined.

DANIEL FREEMAN, BA, PhD, DClInPsy, PHILIPPA A. GARETY, MA, MPhil, PhD, Department of Psychology, Institute of Psychiatry, King's College London; PAUL E. BEBBINGTON, MA, PhD, FRCP, FRCPSych, BENJAMIN SMITH, BSc, DClInPsy, Department of Mental Health Sciences, Royal Free and University College Medical School, London; REBECCA ROLLINSON, BSc, DClInPsy, DAVID FOWLER, BSc, MSc, School of Medicine, Health Policy and Practice, University of East Anglia, Norwich; ELIZABETH KUIPERS, BSc, MSc, PhD, Department of Psychology, KATARZYNA RAY, MA, MSc, Department of Biostatistics and Computing, Institute of Psychiatry, King's College London; GRAHAM DUNN, PhD, Biostatistics Group, Division of Epidemiology and Health Sciences, University of Manchester, Manchester, UK

Correspondence: Dr Daniel Freeman, Department of Psychology, PO Box 77, Institute of Psychiatry, Denmark Hill, London SE5 8AF, UK. E-mail: D.Freeman@iop.kcl.ac.uk

(First received 23 March 2004, final revision 9 September 2004, accepted 10 September 2004)

paranoid thoughts and with angry thoughts and feelings. Attributions that others have negative intentions underlie feelings of anger. We think that in some cases anger may contribute to the attribution of intent in persecutory ideation. However, rather than expressing anger or resentment towards others, individuals may instead ruminate and feel aggrieved owing to timidity or submissiveness. This will maintain a state in which external attributions and anomalous experiences are more likely, thus leading to the persistence of persecutory ideation.

We also found that respondents who felt left out, inferior or less competent in relation to others displayed higher levels of suspiciousness. Birchwood *et al* (2000) reported a connection between social comparison and the experience of hearing voices. We suggest that a lack of social

self-confidence might make people feel vulnerable to attack and hence contribute to the occurrence of paranoia. This is consistent with experimental evidence that interpersonal sensitivity predicts persecutory ideation (Freeman *et al*, 2003). It is of note, however, that in our study the associations of paranoia with many of the variables were of small to medium effect size. This is unsurprising, and is consistent with the view that paranoia is a complex phenomenon likely to arise from a number of social, cognitive and biological factors.

Interventions for paranoid thoughts

Our study has practical implications for clinical interventions in paranoia. Interventions may be more effective if they include recognition of the ubiquity of

suspiciousness; encourage talking about such experiences with others; improve self-esteem; help people in negotiating relationships with others; and encourage detachment and feelings of control over the situation. These are all central components of cognitive-behavioural interventions for psychosis (e.g. Fowler *et al*, 1995; Chadwick *et al*, 1996).

ACKNOWLEDGEMENTS

This work was supported by a programme grant from the Wellcome Trust (no. 062452). We thank the participants from the three universities.

REFERENCES

- Allan, S. & Gilbert, P. (1997) Submissive behaviour and psychopathology. *British Journal of Clinical Psychology*, **36**, 467–488.
- Birchwood, M., Meaden, A., Trower, P., *et al* (2000) The power and omnipotence of voices: subordination and entrapment by voices and significant others. *Psychological Medicine*, **30**, 337–344.
- Birnbaum, M. H. (2001) *Introduction to Behavioural Research on the Internet*. Englewood Cliffs, NJ: Prentice-Hall.
- Chadwick, P. D. J., Birchwood, M. J. & Trower, P. (1996) *Cognitive Therapy for Delusions, Voices and Paranoia*. Chichester: Wiley.
- Fenigstein, A. & Venable, P. A. (1992) Paranoia and self-consciousness. *Journal of Personality and Social Psychology*, **62**, 129–138.
- Folkman, S. & Lazarus, R. S. (1980) An analysis of coping in a middle-aged community sample. *Journal of Health and Social Behaviour*, **21**, 219–239.
- Fowler, D., Garety, P. A. & Kuipers, L. (1995) *Cognitive Behaviour Therapy for Psychosis: Theory and Practice*. Chichester: Wiley.
- Freeman, D. & Garety, P. A. (2004) *Paranoia: The Psychology of Persecutory Delusions*. Hove: Psychology Press.
- Freeman, D., Garety, P. A., Kuipers, E., *et al* (2002) A cognitive model of persecutory delusions. *British Journal of Clinical Psychology*, **41**, 331–347.
- Freeman, D., Slater, M., Bebbington, P. E., *et al* (2003) Can virtual reality be used to investigate persecutory ideation? *Journal of Nervous and Mental Disease*, **191**, 509–514.
- Freeman, D., Garety, P. A., Fowler, D., *et al* (2004) Why do people with delusions fail to choose more realistic explanations for their experiences? An empirical investigation. *Journal of Consulting and Clinical Psychology*, **72**, 671–680.
- Garety, P. A., Kuipers, E., Fowler, D., *et al* (2001) A cognitive model of the positive symptoms of psychosis. *Psychological Medicine*, **31**, 189–195.
- Gilbert, P. & Allan, S. (1994) Assertiveness, submissive behaviour and social comparison. *British Journal of Clinical Psychology*, **33**, 295–306.
- Johns, L. C., Cannon, M., Singleton, N., *et al* (2004) Prevalence and correlates of self-reported psychotic symptoms in the British population. *British Journal of Psychiatry*, **185**, 298–305.
- Joseph, S., Yule, W., Williams, R., *et al* (1994) Correlates of post-traumatic stress at 30 months: the Herald of Free Enterprise disaster. *Behaviour Research and Therapy*, **32**, 521–524.
- Joseph, S., Dalgleish, T., Williams, R., *et al* (1997) Attitudes towards emotional expression and post-traumatic stress in survivors of the Herald of Free Enterprise disaster. *British Journal of Clinical Psychology*, **36**, 133–138.
- Melzer, D., Tom, B. D. M., Brugha, T. S., *et al* (2002) Common mental disorder symptom counts in populations: are there distinct case groups above epidemiological cut-offs? *Psychological Medicine*, **32**, 1195–1201.
- Rachman, S. & de Silva, P. (1978) Abnormal and normal obsessions. *Behaviour Research and Therapy*, **16**, 233–248.
- Roger, D., Jarvis, G. & Najarian, B. (1993) Detachment and coping: the construction and validation of a new scale for measuring coping strategies. *Personality and Individual Differences*, **15**, 619–626.
- SPSS (2001) *SPSS Base 11.0.1 User's Guide*. Chicago, IL: SPSS.
- Sturt, E. (1981) Hierarchical patterns in the distribution of psychiatric symptoms. *Psychological Medicine*, **11**, 783–794.
- Van Os, J. & Verdoux, H. (2003) Diagnosis and classification of schizophrenia: categories versus dimensions, distributions versus disease. In *The Epidemiology of Schizophrenia* (eds R. M. Murray, P. B. Jones, E. Susser, *et al*), pp. 364–410. Cambridge: Cambridge University Press.

BJPpsych

The British Journal of Psychiatry

Psychological investigation of the structure of paranoia in a non-clinical population

Daniel Freeman, Philippa A. Garety, Paul E. Bebbington, Benjamin Smith, Rebecca Rollinson, David Fowler, Elizabeth Kuipers, Katarzyna Ray and Graham Dunn
BJP 2005, 186:427-435.

Access the most recent version at DOI: [10.1192/bjp.186.5.427](https://doi.org/10.1192/bjp.186.5.427)

References

This article cites 16 articles, 1 of which you can access for free at:
<http://bjp.rcpsych.org/content/186/5/427#BIBL>

Reprints/ permissions

To obtain reprints or permission to reproduce material from this paper, please write to permissions@rcpsych.ac.uk

You can respond to this article at

[/letters/submit/bjprcpsych;186/5/427](https://bjp.rcpsych.org/letters/submit/bjprcpsych;186/5/427)

Downloaded from

<http://bjp.rcpsych.org/> on August 19, 2017
Published by The Royal College of Psychiatrists
