

Review article

Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis

Mary Leamy,* Victoria Bird,* Clair Le Boutillier, Julie Williams and Mike Slade

Background

No systematic review and narrative synthesis on personal recovery in mental illness has been undertaken.

Aims

To synthesise published descriptions and models of personal recovery into an empirically based conceptual framework.

Method

Systematic review and modified narrative synthesis.

Results

Out of 5208 papers that were identified and 366 that were reviewed, a total of 97 papers were included in this review. The emergent conceptual framework consists of: (a) 13 characteristics of the recovery journey; (b) five recovery processes comprising: connectedness; hope and optimism about the future; identity; meaning in life; and empowerment

(giving the acronym CHIME); and (c) recovery stage descriptions which mapped onto the transtheoretical model of change. Studies that focused on recovery for individuals of Black and minority ethnic (BME) origin showed a greater emphasis on spirituality and stigma and also identified two additional themes: culturally specific facilitating factors and collectivist notions of recovery.

Conclusions

The conceptual framework is a theoretically defensible and robust synthesis of people's experiences of recovery in mental illness. This provides an empirical basis for future recovery-oriented research and practice.

Declaration of interest

None.

Personal recovery has been defined as 'a deeply personal, unique process of changing one's attitudes, values, feelings, goals, skills and/or roles . . . a way of living a satisfying, hopeful and contributing life even with the limitations caused by illness'.¹ A recovery orientation is mental health policy in most Anglophone countries. For example, the mental health plan for England 2009–2019 has the 'expectation that services to treat and care for people with mental health problems will be . . . based on the best available evidence and focused on recovery, as defined in discussion with the service user'.² The implications of a recovery orientation for working practice are unclear, and guidelines for developing recovery-oriented services are only recently becoming available.^{3,4} Comprehensive reviews of the recovery literature have concluded that there is a need for conceptual clarity on recovery.^{5,6} Current approaches to understanding personal recovery are primarily based on qualitative research⁷ or consensus methods.⁸ No systematic review and synthesis of personal recovery in mental illness has been undertaken.

The aims of this study were (a) to undertake the first systematic review of the available literature on personal recovery and (b) to use a modified narrative synthesis to develop a new conceptual framework for recovery. A conceptual framework, defined as 'a network, or a plane, of interlinked concepts that together provide a comprehensive understanding of a phenomenon or phenomena',⁹ provides an empirical basis for future recovery-oriented research and practice.

illness. A conceptualisation of recovery was defined as either a visual or narrative model of recovery, or themes of recovery, which emerged from a synthesis of secondary data or an analysis of primary data. Inclusion criteria for studies were:

- contains a conceptualisation of personal recovery from which a succinct summary could be extracted;
- presented an original model or framework of recovery;
- was based on either secondary research synthesising the available literature or primary research involving quantitative or qualitative data based on at least three participants;
- was available in printed or downloadable form;
- was available in English.

Exclusion criteria were:

- studies solely focusing on clinical recovery⁴ (i.e. using a predefined and invariant 'getting back to normal' definition of recovery through symptom remission and restoration of functioning);
- studies involving modelling of predictors of clinical recovery;
- studies defining remission criteria or recovery from substance misuse, addiction or eating disorders;
- dissertations and doctoral theses (because of availability).

Search strategy and data sources

Three search strategies were used to identify relevant studies: electronic database searching, hand-searching and web-based searching.

Electronic database searching

Twelve bibliographic databases were initially searched using three different interfaces: Applied and Complimentary Medicine

Method

Eligibility criteria

The review sought to identify papers that explicitly described or developed a conceptualisation of personal recovery from mental

*Joint first authors.

Database (AMED); British Nursing Index; EMBASE; MEDLINE; PsycINFO; Social Science Policy (accessed via OVID SP); CINAHL; International Bibliography of Social Science (accessed via EBSCOhost); Applied Social Science Index and Abstracts (ASSIA); British Humanities Index; sociological abstracts; and Social Services abstracts (accessed via CSA Illumina). All databases were searched from inception to September 2009 using the following terms identified from the title, abstract, keywords or medical subject headings: ('mental health' OR 'mental illness' OR 'mental disorder' OR 'mental disease' OR 'mental problem') AND 'recover' AND ('theor' OR 'framework' OR 'model' OR 'dimension' OR 'paradigm' OR 'concept'). The search was adapted for the individual databases and interfaces as needed. For example, CSA Illumina only allows the combination of three 'units' each made up of three search terms at any one time, for example ('mental health' OR 'mental illness*' OR 'mental disorder') AND 'recover*' AND ('theor*' OR 'framework' OR 'concept'). As a sensitivity check, ten papers were identified by the research team as highly influential, based on number of times cited and credibility of the authors (included papers 3, 9, 10, 19, 29, 34, 35, 40, 68 and 75 in online Table DS1). These papers were assessed for additional terms, subject headings and key words, with the aim of identifying relevant papers not retrieved using the original search strategy. This led to the use of the following additional search terms: ('psychol\$ health' OR 'psychol\$ illness' OR 'psychol\$ disorder' OR 'psychol\$ problem' OR 'psychiatr\$ health' OR 'psychiatr\$ illness' OR 'psychiatr\$ disorder' OR 'psychiatr\$ problem') AND 'recovers' AND ('themes' OR 'stages' OR 'processes'). Duplicate articles were removed within the original database interfaces using Reference Manager Software Version 11 for Windows.

Hand-searching

The tables of contents of journals which published key articles (*Psychiatric Rehabilitation Journal*, *British Journal of Psychiatry* and *American Journal of Psychiatry*) and recent literature reviews of recovery (included papers 4, 37 and 89 in online Table DS1) were hand-searched.

Web-based searching

Web-based resources were identified by internet searches using Google and Google Scholar and through searching specific recovery-oriented websites (Scottish Recovery Network: www.scottishrecovery.net; Boston University Repository of Recovery Resources: www.bu.edu/cpr/repository/index.html; Recovery Devon: www.recoverydevon.co.uk; and Social Perspectives Network: www.spn.org.uk).

Data extraction and quality assessment

One rater (V.B.) extracted data and assessed the eligibility criteria for all retrieved papers, with a random subsample of 88 papers independently rated by a second rater (J.W. or C.L.B.). Disagreements between raters were resolved by a third rater (M.L.). Acceptable concordance was predefined as agreement on at least 90% of ratings. A concordance of 91% was achieved. Data were extracted and tabulated for all papers rated as eligible for the review.

Included qualitative papers were initially quality assessed by three raters (V.B., J.W. and C.L.B.) using the RATS (relevance, appropriateness, transparency, soundness) qualitative research review guidelines.¹⁰ The RATS scale comprises 25 questions about the relevance of the study question, appropriateness of qualitative method, transparency of procedures, and soundness of interpretive approach. In order to make judgements about quality of papers, we dichotomised each question to yes (1 point) or no

(0 points), giving a scale ranging from 0 (poor quality) to 25 (high quality). A random subsample of ten qualitative studies were independently rated using the RATS guidelines by a second rater (M.L.). The mean score from rating 1 was 14.8 and from rating 2 was 15.1, with a mean difference in ratings of 0.3 indicating acceptable concordance. The Effective Public Health Practice Project (EPHPP)¹¹ quality assessment tool for quantitative studies was used to rate the two quantitative studies. Independent ratings were made by two reviewers (V.B. and M.L.) of Ellis & King¹² and Resnick *et al.*,¹³ who agreed on rating both papers as moderate.

Data analysis

The conceptual framework was developed using a modified narrative synthesis approach.¹⁴ The three stages of the narrative synthesis comprised: (1) developing a preliminary synthesis; (2) exploring relationships within and between studies; and (3) assessing the robustness of the synthesis. For clarity, the development of the conceptual framework (Stages 1 and 3) is presented in the Results before the subgroup comparison (Stage 2).

Stage 1: developing a preliminary synthesis

A preliminary synthesis was developed using tabulation, translating data through thematic analysis of good-quality primary data, and vote counting of emergent themes. For each included paper, the following data were extracted and tabulated: type of paper, methodological approach, participant information and inclusion criteria, study location, and summary of main study findings. An initial coding framework was developed and used to thematically analyse a subsample of qualitative research studies with the highest RATS quality rating (i.e. RATS score of 15 or above), using NVIVO QSR International qualitative analysis software (Version 8) for Windows. The main overarching themes and related subthemes occurring across the tabulated data were identified, using inductive, open coding techniques. Additional codes were created by all analysts where needed and these new codes were regularly merged with the NVIVO master copy, and then this copy was shared with other analysts, so all new codes were applied to the entire subsample.

Finally, once the themes had been created, vote counting was used to identify the frequency with which themes appeared in all of the 97 included papers. The vote count for each category comprised the number of papers mentioning either the category itself or a subordinate category. On completion of the thematic analysis and vote counting, the draft conceptual framework was discussed and refined by all authors. Some new categories were created, and others were subsumed within existing categories, given less prominence or deleted. This process produced the preliminary conceptual framework.

Stage 2: exploring relationships within and between studies

Papers were identified from the full review which reported data from people from Black and minority ethnic (BME) backgrounds. These papers were thematically analysed separately, and the emergent themes compared with the preliminary conceptual framework. The thematic analysis utilised a more fine-grained approach, in which a second analyst (V.B.) went through the papers in a detailed and line-by-line manner. The aim of the subgroup analysis was to specifically identify any additional themes as well as any difference in emphasis placed on areas of the preliminary framework. Thus, our purpose was to identify areas of different emphasis in this subgroup of studies, not to perform a validity check.

Stage 3: assessing robustness of the synthesis

Two approaches were used to assess the robustness of the synthesis. First, qualitative studies which were rated as moderate quality on the RATS scale (i.e. RATS score of 14) were thematically analysed until category saturation was achieved. The resulting themes were then compared with the preliminary conceptual framework developed in Stages 1 and 2. Second, the preliminary conceptual framework was sent to an expert consultation panel. The panel comprised 54 advisory committee members of the REFOCUS Programme (see www.researchintorecovery.com for further details) who had academic, clinical or personal expertise about recovery. They were asked to comment on the positioning of concepts within different hierarchical levels of the conceptual framework, identify any important areas of recovery which they felt had been omitted and make any general observations. The preliminary conceptual framework was modified in response to these comments, to produce the final conceptual framework.

Results

The flow diagram for the 97 included papers is shown in Fig. 1 and online Table DS1 lists those papers that were included.

The 97 papers comprised qualitative studies ($n=37$), narrative literature reviews ($n=20$), book chapters ($n=7$), consultation documents reporting the use of consensus methods ($n=5$), opinion pieces or editorials ($n=5$), quantitative studies ($n=2$), combining of a narrative literature review with personal opinion or where there is insufficient information on method for a judgement to be made ($n=11$), and elaborations of other identified papers ($n=10$). In summary, 87 distinct studies were identified. The ten elaborating papers included in the thematic analysis but not in the vote counting were papers 11, 15, 16, 19, 26, 48, 50, 53, 71 and 73 in online Table DS1.

The 97 papers described studies conducted in 13 countries, including the USA ($n=50$), the UK ($n=20$), Australia ($n=8$) and Canada ($n=6$). Participants were recruited from a range of settings, including community mental health teams and facilities, self-help groups, consumer-operated mental health services and supported housing facilities. The majority of studies used inclusion criteria that covered any diagnosis of severe mental illness. A few studies only included participants who had been diagnosed with a specific mental illness (e.g. schizophrenia, depression). The sample sizes in qualitative data papers ranged from 4 to 90 participants, with a mean sample size of 27. The

sample sizes in the two quantitative papers were 19¹² and 1076.¹³ The former was a pilot study of 15 service users with experience of psychotic illness and 4 case managers using the Recovery Interventions Questionnaire, carried out in Australia. The latter study analysed data from two sources, the Schizophrenia Patient Outcomes Research Team (PORT) client survey, which examined usual care in a random sample of people with schizophrenia in two US states, and an extension to this survey which provided a comparison group.

There were various approaches to determining the stage of recovery of participants. Most studies rated stage of recovery using criteria such as: the person defined themselves as 'being in recovery'; not hospitalised during the previous 12 months; relatively well and symptom free; providing peer support to others; or working or living in semi-independent settings. Only a few studies specifically used professional opinion – clinical judgement or scores on clinical assessments – about whether people had recovered.

The mean RATS score for the 36 qualitative studies was 14.9 (range 8–20). One qualitative study was not rated using the RATS guidelines because there was insufficient information on methodology within this paper. A RATS score of 15 or above, indicating high quality, was obtained by 16 papers and used to develop a preliminary synthesis. A RATS score of 14, indicating moderate quality, was obtained by five papers. Independent ratings were made of the two quantitative papers, Ellis & King¹² and Resnick *et al.*,¹³ which were rated as moderate by two reviewers (V.B. and M.L.). Given this quality assessment, no greater weight was put on the quantitative studies in developing the category structure.

Conceptual framework for personal recovery

A preliminary conceptual framework was developed, which comprised five superordinate categories: values of recovery, beliefs about recovery, recovery-promoting attitudes of staff, constituent processes of recovery, and stages of recovery.

The robustness of the synthesis underpinning the preliminary conceptual framework was assessed in two steps: by re-analysing a subsample of qualitative studies and through expert consultation.

Subsample re-analysis

In addition to the higher-quality qualitative studies analysed in the preliminary synthesis stage, an additional five moderate-quality




Fig. 1 Flow chart to show assessment of eligibility of identified studies.

(RATS score of 14) qualitative studies were analysed, which confirmed that category saturation had been achieved, indicating that the categories are robust.

Expert consultation

A response was received from 23 (43%) of the 54 consulted experts with international and national academic, clinical, and/or personal expertise and experiences of recovery, who are advisory committee members of the REFOCUS programme into recovery. Responses were themed under the following headings: conceptual (dangers of reductionism, separating processes from stages, confusing critical impetus for behaviours with actual behaviour, limitations of stage models); structural (complete omissions, lack of emphasis or overemphasis on specific areas of recovery); language (too technical); and bias (potential geographical bias). In response to this consultation, the preliminary conceptual framework was simplified, so the final conceptual framework now has three rather than five superordinate categories. Some subcategories were repositioned within recovery processes, and some category headings changed. Some responses identified areas of omission, such as the role of past trauma, hurt and physical health in recovery. However, no alteration was made to the conceptual framework as these did not emerge from the thematic analysis. Other points regarding the strengths and limitations of the framework are addressed in the Discussion. Overall, the expert consultation process provided a validity check on content of conceptual framework, although we were careful not to make radical changes which would have been unjustified, given the weight of evidence provided from preliminary analysis of the included papers.

The final conceptual framework comprises three interlinked, superordinate categories: characteristics of the recovery journey; recovery processes; and recovery stages.

Characteristics of the recovery journey were identified in all 87 studies, and vote-counting was used to indicate their frequency (Table 1).

The categories of recovery processes and their vote counts, indicating frequency of the process being identified, for the two highest category levels are shown in Table 2.

The full description of recovery processes categories and the vote counting results are shown in online Table DS2.

| Characteristics | Number (%) of 87 studies identifying the characteristics |
|---|--|
| Recovery is an active process | 44 (50) |
| Individual and unique process | 25 (29) |
| Non-linear process | 21 (24) |
| Recovery as a journey | 17 (20) |
| Recovery as stages or phases | 15 (17) |
| Recovery as a struggle | 14 (16) |
| Multidimensional process | 13 (15) |
| Recovery is a gradual process | 13 (15) |
| Recovery as a life-changing experience | 11 (13) |
| Recovery without cure | 9 (10) |
| Recovery is aided by supportive and healing environment | 6 (7) |
| Recovery can occur without professional intervention | 6 (7) |
| Trial and error process | 6 (7) |

Fifteen studies developed recovery stage models. The studies were organised using the transtheoretical model of change,¹⁵ as shown in Table 3.

Recovery in individuals of BME origin

As part of Stage 2 of the narrative synthesis process, six studies of recovery from the perspective of individuals of BME origin were identified within the 87 studies. These six studies were re-analysed by a second analyst (V.B.), using a more fine-grained, line-by-line approach to thematic analysis. These comprised a survey of 50 recipients of a community development project in Scotland,¹⁶ a qualitative interview study of African Americans,¹⁷ a narrative literature review,¹⁸ a qualitative study of 40 Maori and non-Maori New Zealanders,¹⁹ a pilot study to test whether the Recovery Star measure was applicable to Black and Asian ethnic minority populations²⁰ and a mixed method study of 91 males from African–Caribbean backgrounds.²¹ These papers provide some preliminary insights into a small number of distinct ethnic minority perspectives, which do not represent a culturally homogeneous group, although some similarities in experience can be observed. Although these six papers were included in the vote-counting process, four of the six BME papers^{16–18,20} were not used in the first-stage thematic analysis. The line-by-line secondary analysis allowed us to explore in greater detail any differences in emphasis and additional themes present in these papers.

The main finding of the subgroup analysis indicated that there was substantial similarity between studies focusing on ethnic minority communities and those focusing on ethnic majority populations. All of the themes of the conceptual framework were present in all six of the BME papers. Despite this overall similarity, there was a greater emphasis in the BME papers on two areas in the recovery processes: spirituality and stigma; and two additional

| Recovery processes | Number (%) of 87 studies identifying the process |
|--|--|
| Category 1: Connectedness | 75 (86) |
| Peer support and support groups | 39 (45) |
| Relationships | 33 (38) |
| Support from others | 53 (61) |
| Being part of the community | 35 (40) |
| Category 2: Hope and optimism about the future | 69 (79) |
| Belief in possibility of recovery | 30 (34) |
| Motivation to change | 15 (17) |
| Hope-inspiring relationships | 12 (14) |
| Positive thinking and valuing success | 10 (11) |
| Having dreams and aspirations | 7 (8) |
| Category 3: Identity | 65 (75) |
| Dimensions of identity | 8 (9) |
| Rebuilding/redefining positive sense of identity | 57 (66) |
| Overcoming stigma | 40 (46) |
| Category 4: Meaning in life | 59 (66) |
| Meaning of mental illness experiences | 30 (34) |
| Spirituality | 6 (41) |
| Quality of life | 57 (65) |
| Meaningful life and social roles | 40 (46) |
| Meaningful life and social goals | 15 (17) |
| Rebuilding life | 19 (22) |
| Category 5: Empowerment | 79 (91) |
| Personal responsibility | 79 (91) |
| Control over life | 78 (90) |
| Focusing upon strengths | 14 (16) |

Table 3 Recovery stages mapped onto the transtheoretical model of change

| Table DS1 | | | | | |
|--------------|-------------------------------|---|--|--|--|
| study number | Precontemplation | Contemplation | Preparation | Action | Maintenance and growth |
| 32 | | Novitiate recovery – struggling with disability | | Semi-recovery – living with disability | Full recovery – living beyond disability |
| 73 | Stuck | Accepting help | Believing | Learning | Self-reliant |
| 3 | Descent into hell | Igniting a spark of hope | Developing insight/ activating instinct to fight back | Discovering keys to well-being | Maintaining equilibrium between internal and external forces |
| 44 | Demoralisation | | Developing and establishing independence | | Efforts towards community integration |
| 36 | Occupational dependence | | Supported occupational performance | Active engagement in meaningful occupations | Successful occupational performance |
| 14 | Dependent/unaware | Dependent/aware | | Independent/aware | Interdependent/aware |
| 29 | Moratorium | Awareness | Preparation | Rebuilding | Growth |
| 78 | | Glimpses of recovery | Turning points | Road to recovery | |
| 61 | | Reawakening of hope after despair | No longer viewing self as primarily person with psychiatric disorder | Moving from withdrawal to engagement | Active coping rather than passive adjustment |
| 40 | Overwhelmed by the disability | | Struggling with the disability | Living with the disability | Living beyond the disability |
| 35 | Initiating recovery | | | Regaining what was lost/moving forward | Improving quality of life |
| 59 | Crisis (recuperation) | | Decision (rebuilding independence) | Awakening (building healthy interdependence) | |
| 43 | | Turning point | Determination | | Self-esteem |

categories: culture-specific factors and collectivist notions of recovery.

In relation to spirituality, being part of a faith community and having a religious affiliation was seen as an important component of an individual's recovery. People from ethnic minority groups more often described spirituality in terms of religion and a belief in God as a higher power, whereas participants in the non-BME studies tended to conceptualise spirituality as encompassing a wider range of beliefs and activities.

In relation to stigma, BME studies emphasised the stigma associated with race, culture and ethnicity, in addition to the stigma associated with having a mental illness. Furthermore, being an individual from a minority ethnic group seemed to accentuate the stigma of mental illness, as the person often viewed themselves as belonging to multiple stigmatised and disadvantaged groups. Individuals from ethnic minority groups saw themselves as recovering from racial discrimination, stigma and violence, and not just from a period of mental illness.

The new category of culture-specific factors included the use of traditional therapies and faith healers, and belonging to a particular cultural group or community. Finally, collectivist notions of recovery were emphasised as both positive and negative factors. Many individuals discussed the hope and support they received from their collectivist identity, but for others the community added to the pressures of mental illness. This was particularly true where communities lacked information and awareness regarding mental illness. Furthermore, the negative impact of the community was felt not only at the level of the individual, but also at the collectivist level, with the whole family being adversely affected by stigma.

Discussion

This is the first systematic review and narrative synthesis of personal recovery. A conceptual framework was developed using a narrative synthesis which identified three superordinate

categories: characteristics of the recovery journey, recovery processes and recovery stages. For each superordinate category, key dimensions were synthesised. The recovery processes that have the most proximal relevance to clinical research and practice are: connectedness; hope and optimism about the future; identity; meaning in life; and empowerment (giving the acronym CHIME). The robustness of the category structure was enhanced by the systematic nature of the review, the quality assessment of included studies, the category saturation reached in the analysis, and the content validity of the expert consultation. Heterogeneity between studies was explored descriptively. A subgroup comparison between the experiences of recovery from the perspective of individuals of BME origin identified similar themes, with a greater emphasis on spirituality and stigma, and two additional themes: culture-specific factors, and collectivist notions of recovery.

Implications for research and practice

Key knowledge gaps have been identified as the need for clarity about the underpinning philosophy of recovery,²² better understanding of the stages and processes of recovery,⁵ and valid measurement tools.²³ This study can inform each of these gaps.

Recovery has been conceptualised as a vision, a philosophy, a process, an attitude, a life orientation, an outcome and a set of outcomes.⁵ This has led to the concern that 'its scope can make a cow-catcher on the front of a road train look discriminating'.²⁴ An empirically based conceptual framework can bring some order to this potential chaos. Characteristics of the recovery journey provide conceptual clarity about the philosophy. Recovery processes can be understood as measurable dimensions of change, which typically occur during recovery and provide a taxonomy of recovery outcomes.²⁵ Finally, recovery stages provide a framework for guiding stage-specific clinical interventions and evaluation strategies.

The framework contributes to the understanding about stages and processes of recovery in two ways. First, it allows available

evidence to be more easily identified. A recovery orientation has overlap with the literature on well-being,²⁶ positive psychology²⁷ and self-management,²⁸ and systematic reviewing is hampered by the absence of relevant MeSH (Medical Sub-Headings) headings relating to recovery concepts. The coding framework provides keywords for use when undertaking secondary research, and the identification of related terms provides a taxonomy which will be useable in reviews.

Second, the framework provides a structure around which research and clinical efforts can be oriented. The relative contribution of each recovery process, investigating interventions which can support these processes, and the synchrony between recovery processes and stages are all testable research questions. For clinical practice, the CHIME recovery processes support reflective practice. If the goal of mental health professionals is to support recovery then one possible way forward is for each working practice to be evaluated in relation to its impact on these processes. This has the potential to contribute to current debates about recovery and, for example, assertive outreach,²⁹ risk³⁰ and community psychiatry.³¹

Finally, the conceptual framework can contribute to the development of measures of personal recovery. Compendia of existing measures have been developed,^{32,33} showing that the conceptual basis of measures is diverse. The conceptual framework provides a foundation for developing standardised recovery measures, and is the basis for a new measure currently being developed by the authors to evaluate the contribution of mental health services to an individual's recovery. The challenge will then be to incorporate a focus on recovery outcomes and associated concepts such as well-being²⁷ into routine clinical practice.³⁴

Limitations

The study has three methodological and two conceptual limitations. The first methodological limitation is that the narrative synthesis approach was modified, and could have been widened. For example, the exploration in Stage 2 of relationships between studies could have considered the subgroup of studies which had higher levels of consumer involvement in their design, but it proved impossible to reliably rate identified studies in this dimension. The second technical limitation is that the emergent categories were only one way of grouping the findings, and the categories changed as a result of expert consultation. In particular, the three superordinate categories are not separate, since processes clearly occur within the identified stages, and the characteristics of recovery describe an overall movement through stages of recovery. Our categorisation brings structure, but a replication study may not arrive at the same overall thematic structure. The final technical limitation is that analysis synthesised the interpretation of the primary data in each paper rather than considering the primary data directly. Future research could compare papers generated by different stakeholder groups, such as consumer researchers, clinical researchers and policy makers.

The first conceptual limitation is that this review, although synthesising the current literature on personal recovery, should not be seen as definitive. A key scientific challenge is that the philosophy of recovery gives primacy to individual experience and meaning ('idiographic' knowledge), whereas mental health systems and current dominant scientific paradigms give prominence to group-level aggregated data ('nomothetic' knowledge).⁴ The practical impact is that current recovery research is primarily focused at the bottom of the hierarchy of evidence.³⁵ This was our finding, with qualitative, case study and expert opinion methodologies dominating. A motivator for the current study was to provide evidence of the form viewed as

high quality within the current scientific paradigm, but several of our expert consultants highlighted the dangers of closing down discourse. Since recovery is individual, idiosyncratic and complex, this review is not intended to be a rigid model of what recovery 'is'. Rather, it is better understood as a resource to inform future research and clinical practice. The second conceptual limitation relates to the subgroup analysis looking at papers focusing on non-majority populations. Owing to a lack of research, it was not possible to look at the experience and perspectives of individuals from different ethnic minority groups. Therefore, the BME subgroup represents a heterogeneous and incredibly diverse set of populations. However, it was felt that all the populations included in these papers shared a common experience of belonging to an ethnic minority group, and that this experience may have important implications for the meaning of personal recovery, and for the experience of mental health services in general. The lack of data, coupled with the areas of difference found in the present review, highlights a need for further work to be conducted with people from ethnic minority communities.

Future research

This systematic review and narrative synthesis has highlighted the dominance of recovery literature emanating from the USA. Culturally, the USA neglects character strengths such as patience and tolerance,³⁶ and favours individualistic over collectivist understandings of identity. Although there were very few studies which looked at recovery experiences of individuals from BME backgrounds, the subsample of BME studies indicated that there are important differences in emphasis. There is a need for research involving more diverse samples of people from different ethnic and cultural backgrounds, at differing stages of recovery and experiencing different types of mental illness.

The complexity of personal recovery requires a range of theoretical inquiry positions. This review focused on research into first-person accounts of recovery, where individual meanings of recovery have dominated. This has led to a framework which may underemphasise the importance of the wider socio-environmental context, including important aspects such as stigma and discrimination. Viewing recovery within an ecological framework, as suggested by Onken and colleagues,³⁵ encompasses an individual's life context (characteristics of the individual such as hope and identity) as well as environmental factors (such as opportunities for employment and community integration) and the interaction between the two (such as choice). A more complete understanding of recovery requires greater attention to all these levels of understanding, for instance, how power is related to characteristics of individuals or groups (e.g. race and culture), how clinicians and patients interact at different stages of recovery and how these interactions change over time. There is also a need for future research to increase our understanding of how subtle micro-processes of recovery are operating, such as how hope is reawakened and sustained.

Supporting recovery processes may be the future mental health research priority. The 13 dimensions identified as characteristics of the recovery journey capture much of the experience and complexities of recovery, and further research may not have a high scientific pay-off. Similarly, although the recovery stages could be mapped onto the transtheoretical model of change,¹⁵ there was little consensus about the number of recovery phases. It may therefore be more helpful to undertake evaluative research addressing specific service-level questions (such as whether people using a service are making recovery gains over time³⁷ or in different service settings³⁸), rather than further studies seeking conceptual clarity. Overall, the emergent priority

is the development and evaluation of interventions to support the five CHIME recovery processes. The subordinate categories point to the need for a greater emphasis on assessment of strengths and support for self-narrative development, promoting the role of mental health systems in developing inclusive communities enabling access to peer support as well as providing retreats, and clinical interaction styles which promote empowerment and self-management. The CHIME categories are potential clinical end-points for interventions, in contrast to the current dominance of clinical recovery end-points such as symptomatology or hospitalisation rates. They also provide a framework for empirical investigation of the relationship between recovery outcomes, using methodologies developed in relation to clinical outcomes.³⁹ This area of enquiry is currently small⁴⁰ but an important priority if potential trade-offs between desirable outcomes are to be identified.⁴¹

Orienting mental health services towards recovery will involve system transformation.⁴² The research challenge is to develop an evidence base which simultaneously helps mental health professionals to support recovery and respects the understanding that recovery is a unique and individual experience rather than something the mental health system does to a person. This conceptual framework for personal recovery, which has been developed through a systematic review and narrative synthesis, provides a useful starting point for meeting this challenge.

Mary Leamy, PhD, **Victoria Bird**, BSc, **Clair Le Bouillier**, MSc, **Julie Williams**, MSc, **Mike Slade**, PhD, PsychD, Health Service and Population Research Department, King's College London, Institute of Psychiatry, UK

Correspondence: Mary Leamy PhD, Programme Co-ordinator, King's College London, Health Service and Population Research Department (Box P029), Institute of Psychiatry, De Crespigny Park, Denmark Hill, London SE5 8AF, UK. Email: mary.leamy@kcl.ac.uk

First received 23 Jun 2010, final revision 21 Apr 2011, accepted 26 May 2011

Funding

This study was funded by a National Institute for Health Research (NIHR) Programme Grant for Applied Research (RP-PG-0707-10040) awarded to the South London and Maudsley NHS Foundation Trust, and in relation to the NIHR Specialist Mental Health Biomedical Research Centre at the Institute of Psychiatry, King's College London. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health.

Acknowledgements

We would like to thank Popay *et al* for giving us the unpublished guidance on narrative synthesis.

References

- Anthony WA. Recovery from mental illness: the guiding vision of the mental health system in the 1990s. *Innovations and Research* 1993; **2**: 17–24.
- HM Government. *New Horizons. A Shared Vision for Mental Health*. Department of Health, 2009.
- Davidson L, Tondora J, Lawless MS, O'Connell M, Rowe M. *A Practical Guide to Recovery-Oriented Practice Tools for Transforming Mental Health Care*. Oxford University Press, 2009.
- Slade M. *Personal Recovery and Mental Illness. A Guide for Mental Health Professionals*. Cambridge University Press, 2009.
- Silverstein SM, Bellack A. A scientific agenda for the concept of recovery as it applies to schizophrenia. *Clin Psychol Rev* 2008; **28**: 1108–24.
- Warner R. Recovery from schizophrenia and the recovery model. *Curr Opin Psychiatry* 2009; **22**: 374–80.
- Davidson L, Schmutte T, Dinzeo T, Andres-Hyman R. Remission and recovery in schizophrenia: practitioner and patient perspectives. *Schizophr Bull* 2008; **34**: 5–8.
- Substance Abuse and Mental Health Services Administration. *National Consensus Statement on Mental Health Recovery*. Department of Health and Human Services, 2006.
- Jabareen Y. Building a conceptual framework: philosophy, definitions, and procedure. *Int J Qual Meth* 2009; **8**: 49–62.
- Clark J. How to peer review a qualitative manuscript. In *Peer Review in Health Sciences (2nd edn)*. (eds F Godlee, T Jefferson): 219–35. BMJ Books, 2003.
- Effective Public Health Practice Project. *Quality Assessment Tool for Quantitative Studies*. EPHPP, 2009 (<http://www.ephpp.ca/tools.html>).
- Ellis G, King R. Recovery focused interventions: perceptions of mental health consumers and their case managers. *AeJAMH* 2003; **2**: 067–076.
- Resnick SG, Fontana A, Lehman A, Rosenheck RA. An empirical conceptualization of the recovery orientation. *Schizophr Res* 2005; **75**: 119–28.
- Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, et al. Guidance on the conduct of narrative synthesis in systematic reviews. Results of an ESRC funded research project (unpublished report). University of Lancaster, 2006.
- Prochaska JO, DiClemente CC. Transtheoretical therapy: toward a more integrative model of change. *Psychol Psychother Theor Res Pract* 1982; **19**: 276–88.
- NHS Scotland. *Finding Strength From Within. Report on Three Local Projects Looking at Mental Health and Recovery with People from some of the Black and Minority Ethnic Communities in Edinburgh*. NHS Scotland, 2008.
- Armour R, Bradshaw W, Roseborough D. African Americans and recovery from severe mental illness. *Soc Work Ment Health* 2009; **7**: 602–22.
- Nicholls V. *Feeding the Flowers: SPN Perspective on Recovery*. Social Perspectives Network, 2007.
- Lapsley H, Nikora LW, Black R. *Kia Mauri Tau! Narratives of Recovery from Disabling Mental Health Problems*. Mental Health Commission, 2002.
- Mental Health Providers Forum. *The Recovery Star Model and Cultural Competency*. Mental Health Providers Forum, 2009.
- Brown M, Essien P, Etim-Ubah P. *Report of the Community Led Research Project Focusing on Male African and African Caribbean Perspectives on Recovery*. Fanon Southside Partnership, 2008.
- Dinniss S. Recovery-oriented mental healthcare. *Br J Psychiatry* 2006; **189**: 384.
- Care Services Improvement Partnership, Royal College of Psychiatrists, Social Care Institute for Excellence. *A Common Purpose: Recovery in Future Mental Health Services*. CSIP, 2007.
- Brunskill D. From patient to service user – in 100 words. *Br J Psychiatry* 2010; **196**: 353.
- Slade M. What outcomes to measure in routine mental health services, and how to assess them: a systematic review. *Aust N Z J Psychiatry* 2002; **36**: 743–53.
- Hanlon P, Carlisle S. What can the science of well-being tell the discipline of psychiatry – and why might psychiatry listen? *Adv Psychiatr Treat* 2008; **14**: 312–9.
- Slade M. Mental illness and well-being: the central importance of positive psychology and recovery approaches. *BMC Health Serv Res* 2010; **10**: 26.
- Sterling EW, von Esenwein SA, Tucker S, Fricks L, Druss BG. Integrating wellness, recovery, and self-management for mental health consumers. *Community Ment Health J* 2010; **46**: 130–8.
- Drake RE, Deegan P. Are assertive community treatment and recovery compatible? *Community Ment Health J* 2008; **44**: 75–7.
- Young AT, Green CA, Estroff SE. New endeavors, risk taking, and personal growth in the recovery process: findings from the STARS study. *Psychiatr Serv* 2008; **59**: 1430–6.
- Rosen A. The community psychiatrist of the future. *Curr Opin Psychiatry* 2006; **19**: 380–8.
- Campbell-Orde T, Chamberlin J, Carpenter J, Leff SH. *Measuring the Promise: A Compendium of Recovery Measures. Vol II*. Human Services Research Institute, 2005.
- National Institute for Mental Health in England. *Outcomes Compendium*. NIMHE, 2008.
- Slade M. Routine outcome assessment in mental health services. *Psychol Med* 2002; **32**: 1339–43.
- Onken SJ, Craig CM, Ridgway P, Ralph RO, Cook JA. An analysis of the definitions and elements of recovery: a review of the literature. *Psychiatr Rehabil J* 2007; **31**: 9–22.
- Henry J. Positive psychology and the development of well-being. In *Well-being: Individual, Community and Societal Perspectives* (eds J Haworth, G Hart): 25–40. Palgrave Macmillan, 2007.

- 37 Miller L, Brown TT, Pilon D, Scheffler RM, Davis M. Patterns of recovery from severe mental illness: a pilot study of outcomes. *Community Ment Health J* 2010; **46**: 177–87.
- 38 Johnson S, Gilbert H, Lloyd-Evans B, Osborn DPJ, Boardman J, Leese M, et al. In-patient and residential alternatives to standard acute psychiatric wards in England. *Br J Psychiatry* 2009; **194**: 456–63.
- 39 Salvi G, Leese M, Slade M. Routine use of mental health outcome assessments: choosing the measure. *Br J Psychiatry* 2005; **186**: 146–52.
- 40 Andresen R, Caputi P, Oades L. Do clinical outcome measures assess consumer-defined recovery? *Psychiatry Res* 2010; **177**: 309–17.
- 41 Slade M, Hayward M. Recovery, psychosis and psychiatry: research is better than rhetoric. *Acta Psychiatr Scand* 2007; **116**: 81–3.
- 42 Shepherd G, Boardman J, Burns M. *Implementing Recovery. A Methodology for Organisation Change*. Sainsbury Centre for Mental Health, 2010.



poems
by
doctors

Street-wise

Wendy French

Somewhere in a North London street near Northwick Park
a retired doctor pins his butterflies, worries about sex.

Elsewhere in the street a siren stops outside
a two-up, two-down where paramedics collect
a woman with a fluttering heart.

Somewhere in an East London street near the London
a young doctor revising for her MRCP dreams of take-aways,

Chicken Biryani, Tarka Dal.
Elsewhere a man dials 999, he doesn't want to die.
Whisky half drunk and paracetamol gone.

Somewhere in a South London street near King's
a woman gazes into the gas fire, thinks about her husband

and the locum who came at 3am, drank tea.
His own father recently dead and his mother who keeps
all doors unlocked for her husband's return.

Elsewhere near UCH a woman's contractions
Increase as she phones her partner, he's not at his desk

so the neighbour drives her as fast as he can and leaves her
to the student on duty who comforts her between groans.
The partner enjoys a light lunch. Pint of pride.

Somewhere in a London street a man comes near to dying.
His car skids sideways on ice. Approaching cars close in.

He thinks after one death there is no other.
A stranger helps him shuffle along
until stillness returns.

In the same street the old doctor remembers climbing
the snow-ridden hills, a bride by his side and he still feels

her bracelets, purse, red felt hat.
And there's his grandson with Down's who loves to touch velvet,
collect stamps and who lives in a home on a West London street

where the Hammersmith closes. Wards full of flu-ridden adults.
Next door to the doctor little Louise in the wheelchair

drinks orange through a straw, cries throughout the night.
The physician turns back to his moth, the Bloodvein,
a splayed sacrifice and sighs.

This poem is from *The Hippocrates Prize 2011*, published by The Hippocrates Prize in association with Top Edge Press.

Chosen by Femi Oyeboode.

Table DS1 Included studies (n=97)

| | Full reference | Country | Method | Quality rating |
|----|---|------------------|--|----------------|
| 1 | Provencher H, Gregg R, Mead S, Mueser K. The role of work in the recovery of persons with psychiatric disabilities, <i>Psychiatr Rehab. J</i> , 2002, 26 (2), 132-144. | USA | Semi-structured individual interviews (n=14 participants with psychiatric disabilities) | 13/25 |
| 2 | Kelly M, Gamble C. Exploring the concept of recovery in schizophrenia, <i>J Psychiatr Ment Health Nurs</i> . 2005 Aug; 12 (4):386. | Unclear | Literature review | Not rated |
| 3 | Noiseux S, Ricard N. Recovery as perceived by people with schizophrenia, family members and health professionals: A grounded theory. <i>Int. J of Nurs. Studies</i> , 2008, 45 (8), 1148-1162 | Canada | Semi-structured interviews and field notes (n=41 people with schizophrenia, family members and health professionals) | 18/25 |
| 4 | Social Care Institute for Excellence, A common purpose: Recovery in future mental health services, 2007. | UK | Literature review | Not rated |
| 5 | Schon UK, Denhov A, Topor A. Social relationships as a decisive factor in recovering from severe mental illness, <i>Int. J of Soc Psychiatry</i> , 2009, 55 (4) 336-347. | Sweden | Interviews (n=58 people who had recovered from serious mental illness) | 13/25 |
| 6 | Smith M. Recovery from severe psychiatric disability: Findings of a qualitative study, <i>Psychiatr Rehab. J</i> , 2000, 24 (2), 149-158 | USA | Semi-structured interviews (n=10 participants with serious mental illness) | 13/25 |
| 7 | Tooth B, Kalyanasundaram V, Glover H, Momenzadah S. Factors consumers identify as important to recovery from schizophrenia, <i>Australasian Psychiatry</i> , 2003, 11 (1), 70-77. | Australia | Focus groups (n=57 people in recovery from schizophrenia) | 12/25 |
| 8 | Libermann R, Kopelowicz A, Ventura J, Gutkind D. Operational criteria and factors related to recovery from schizophrenia, <i>Int. Review of Psychiatry</i> , 2002a. 14 (4), 256-272. | USA | Literature review, focus groups, case vignettes of people recovering from schizophrenia | Not rated |
| 9 | Ramon S, Healy B, Renouf N. Recovery from Mental Illness as an Emergent Concept and Practice in Australia and the UK, <i>Int.. J. Soc Psychiatry</i> , 53 , 108-122. | UK and Australia | Literature review | Not rated |
| 10 | Mancini MA. A qualitative analysis of turning points in the recovery process, <i>American J of Psychiatr Rehab.</i> , 2007, 10 (3), 223-244. | USA | Semi-structured interviews (n=16 participants recovering from serious psychiatric disability) | 13/25 |
| 11 | Mezzina R, Davidson L, Borg M, Marin I, Topor A, Sells D. The social nature of recovery: Discussion and implications for practice, <i>American J of Psychiatr Rehab.</i> 2006, 9(1), 63-80. | Italy and USA | Literature review and conceptual paper | Not rated |
| 12 | Fallot R. Spiritual and religious dimensions of mental illness recovery narratives, <i>New directions for mental health services</i> , 80 , Winter, 1998. | Unclear | Personal narratives and literature review | Not rated |
| 13 | Morse G. On being homeless and mentally ill: A multitude of losses and the possibility of recovery, chapter 16, in Harvey J & Miller E (Eds). <i>Loss and trauma: General and close relationship perspectives</i> . New York, US: Brunner-Routledge, 2000 | Unclear | Personal narratives and literature review | Not rated |
| 14 | <i>Emerging best practices in mental health recovery</i> , National Institute for Mental Health in England., Great Britain. National Health Service, 2004. | UK | Based on Ohio Department of Mental Health work on the meaning and process of recovery. | Not rated |
| 15 | Piat M, Sabetti J, Couture A, Sylvestre J, Provencher H, Botschner J, Stayner D. What does recovery mean for me? Perspectives of Canadian mental health consumers, <i>Psychiatr Rehab. J</i> , 2009, 32 (3), 199-207. | Canada | Qualitative interviews (n= 60 consumers of mental health services) | 18/25 |

| | | | | |
|----|--|-----------|---|----------------------|
| 16 | Davidson L, O'Connell M, Staeheli M, Weingarten R, Tondora J, Evans A. Concepts of recovery in Behavioral health: History, review of the evidence, and critique, in Davidson L, Rowe M, Tondora J, O'Connell M, Lawless M, <i>A practical guide to recovery-oriented practice</i> . Oxford. Oxford University Press, 2009. | USA | Literature review | Not rated |
| 17 | Diamond R. Recovery from a psychiatrist's viewpoint, <i>Postgraduate Medicine</i> , 2006, Sept. Special, 54-62 | Unclear | Literature review | Not rated |
| 18 | Gagne C, White W, Anthony W. Recovery: A common Vision for the fields of mental health and addiction, <i>Psychiatr Rehab. J</i> , 2007, 31 (1), 32-37. | USA | Literature review | Not rated |
| 19 | Davidson L, O'Connell M, Tondora J, Lawless M, Evans A. Recovery in Serious Mental Illness: A New Wine or Just a New Bottle? <i>Professional Psychology: Research and Practice</i> , 36 (5), 480-487 | USA | Literature review and concept map | Not rated |
| 20 | Davidson L, O'Connell M, Staeheli M, Weingarten R, Tondora J, Evans A A model of being in recovery as a foundation for recovery-oriented practice, in Davidson L, Rowe M, Tondora J, O'Connell M, Lawless M <i>A practical guide to recovery-oriented practice</i> . Oxford. Oxford University Press, 2009. | USA | Interviews (n=100 consumers and people who have lived with mental illness) | Not rated |
| 21 | Slade M. 'Recovery-focussed mental health services: The personal recovery framework', in <i>Personal recovery and mental illness: A guide for mental health professionals</i> , Cambridge University Press, 2009. | UK | Literature review | Not rated |
| 22 | Repper J, Perkins R. 'The individual's recovery journey: towards a model for mental health practice, in Repper, J. & Perkins, R. <i>Social inclusion and recovery: a model for mental health practice</i> , Bailliere Tindall, 2003. | UK | Literature review | Not rated |
| 23 | Markowitz FE. Sociological Models of Recovery, chapter 4, in Ralph, R. & Corrigan, P. <i>Recovery in mental illness: Broadening our understanding of wellness</i> . Washington, DC, US: American Psychological Association, 2005. | USA | Literature review | Quality not assessed |
| 24 | Ralph R. Verbal Definitions and Visual Models of Recovery: Focus on the Recovery Model, in Ralph R, Corrigan P. <i>Recovery in Mental illness: Broadening our understanding and wellness</i> , Washington, DC, US: American Psychological Association, 2005. | USA | Literature review | Not rated |
| 25 | Libermann RP, Kopelowicz A. Recovery from schizophrenia: A challenge for the 21st century, <i>Int. Review of Psychiatry</i> , 2002, 14(4) 245-255. | USA | Literature review | Not rated |
| 26 | Libermann R, Kopelowicz A. Open forum. Recovery from schizophrenia: a concept in search of research, <i>Psychiatr Services</i> , 2005, 56 (6), 735-742 | USA | Literature review | Not rated |
| 27 | Whitehorn D, Brown J, Richard J, Rui Q, Kopala L Multiple dimensions of recovery in early psychosis, <i>Int. Review of Psychiatry</i> , 2002, 14(4), 273-283. | Canada | Literature review | Not rated |
| 28 | Ellis G, King R. Recovery focused interventions: Perceptions of mental health consumers and their case managers. <i>Australian e-J for the Advancement of Ment. Health</i> , 2003, 2 (2). | Australia | Literature review and piloting of a consumer and case manager questionnaire | Not rated |
| 29 | Andresen R, Oades L, Caputi P. The experience of recovery from schizophrenia: towards an empirically validated stage model, <i>Australian & New Zealand J of Psychiatry</i> , 2003, 37 (5), 586-594. | Australia | Literature review and qualitative analysis | Not rated |
| 30 | Torrey W, Wyzik, P. The recovery vision as a service improvement guide for community mental health center providers. <i>Community Ment. Health J</i> , 2000, 36 (2):209-216. | USA | Opinion piece | Not rated |
| 31 | Cleary A, Dowling M. The road to recovery, <i>Ment. Health Practice</i> , 2009, 12 (5), 28- | Ireland | Literature review | Not rated |

| | | | | |
|----|--|-------------|--|-----------|
| | 31. | | | |
| 32 | Song L-Y, Shih C-Y. Factors, process and outcome of recovery from psychiatric disability the utility model, <i>Int. J of Social Psychiatry</i> , 2009, 55 (4), 348-360. | Taiwan | Qualitative interviews (n=15 consumers in recovery and their caregivers) | 15/25 |
| 33 | Resnick S, Fontana A, Lehman A, Rosenheck R. An empirical conceptualization of the recovery orientation, <i>Schizophrenia Research</i> , 2005, 75 , 119-128. | USA | Survey on the treatment of schizophrenia (n=1,076) | Not rated |
| 34 | Jacobson N. Experiencing recovery: A dimensional analysis of recovery narratives, <i>Psychiatr Rehab. J</i> 2001 Winter; 24 (3):248-56. | USA | Dimensional analysis of 30 narratives of recovery. | Not rated |
| 35 | Young S, Ensing D. Exploring recovery from the perspective of people with psychiatric disabilities, <i>Psychiatr Rehab. J</i> , 1999, 22 (3), 219-231. | USA | Semi-structured interviews (n= 18 people with psychiatric disabilities) and focus groups (n=2, 11 participants in total) | 15/25 |
| 36 | Merryman M, Riegel S. The recovery process and people with serious mental illness living in the community: An occupational therapy perspective, <i>Occupational Therapy in Ment. Health</i> . 2007, 23 (2), 51-73. | USA | Interviews (n=20 service users) | 16/25 |
| 37 | Ralph R. Recovery, <i>Psychiatr Rehab. Skills</i> , 2000, 4 (3), 480-517. | USA | Literature review | Not rated |
| 38 | Jensen L, Wadkins T. Mental health success stories: finding paths to recovery, <i>Issues in Ment. Health Nurs.</i> , 2007, 28 (4), 325-340. | USA | Semi-structured interviews (n=20 service users) | 13/25 |
| 39 | Schrank B, Slade M. Recovery in psychiatry, <i>Psychiatr Bulletin</i> , 2007, 31 (9), 321-325. | Austria, UK | Literature review | Not rated |
| 40 | Spaniol S, Wewiorski N, Gagne C, Anthony W. The process of recovery from schizophrenia, <i>Int. review of psychiatry</i> , 2002, 14 , 327-336. | USA | Interviews (n=12 consumers, conducted every four to eight months, over a 4 year period) | 16/25 |
| 41 | Mental Health Recovery Study Working group, <i>Mental Health 'Recovery': users and refusers. What do psychiatric survivors in Toronto think about Mental Health Recovery?</i> Wellesley Institute, 2009. | Canada | Community-based participatory research approach., focus groups (n=7) | Not rated |
| 42 | Hopper K. Rethinking social recovery in schizophrenia: what a capabilities approach might offer, <i>Social Science and Medicine</i> , 2007, 65 (5), 868-879. | USA | Literature review | Not rated |
| 43 | Peden A. Recovering in depressed women: research with Peplau's theory. <i>Nurs Sci Q</i> , 1993, 6 (3), 140-146 | USA | Semi-structured interviews (n= 7 participants recovering from depression) | 14/25 |
| 44 | Bradshaw W, Armour M, Roseborough D. Finding a place in the World: The experience of Recovery from Severe Mental Illness, <i>Qual. Social Work</i> , 2007, 6 (1), 27-47. Sage Publications, UK. | USA | Semi-structured interviews (n= 45 with severe mental illness, conducted over 3 years) | 18/25 |
| 45 | Sung K, Kim S, Puskar K, Kim E. Comparing Life Experiences of College Students with Differing Courses of Schizophrenia in Korea: Case Studies Perspectives in <i>Psychiatric Care</i> , 2006, 42 (2), 82-94. | South Korea | In-depth interviews (n= 8 people diagnosed with schizophrenia) | 17/25 |
| 46 | NHS Scotland, Finding strength from within, Report on three local projects looking at mental health and recovery with people from some of black and minority ethnic communities in Edinburgh, 2008. | Scotland | Exploratory community development project (n= 50 people from BME communities with personal experience of recovery) | Not rated |
| 47 | Ajayi S, Billsborough J, Bowyer T, Brown P, Hicks A, Larsen J, Mailey P, Sayers R, Smith R. Getting back into the world: Reflections on lived experiences of recovery, <i>Rethink recovery series: 2.</i> , 2009. | UK | Interviews (n=48 people with personal experience of mental illness) | 18/25 |
| 48 | Connecticut Department of Mental Health Addiction Services: Proposed model of mental health recovery and recovery-oriented services, in Davidson L, Rowe M, Tondora J, O'Connell M, Lawless M, <i>A practical guide to recovery-oriented practice</i> . Oxford. Oxford University Press, 2009. | USA | Position paper | Not rated |

| | | | | |
|----|--|-----------|---|------------|
| 49 | Mancini A. Self-determination theory: A framework for the recovery paradigm, <i>Adv. in Psychiatr Treatment</i> .2008, 14 (5),358-365. | USA | Literature review | Not rated |
| 50 | Armour M, Bradshaw W. Roseborough D. African Americans and recovery from severe mental illness, <i>Social Work in Ment. Health</i> , 2009, 7 (6), 602-622. | USA | Semi-structured interviews (n=9 African-American with serious and persistent mental illness, conducted with each participant 3 times) | 11/25 |
| 51 | Davidson L, Andres-Hyman R, Bedregal L, Tondora J, Fry J, Kirk T. From 'Double trouble to Dual recovery': Integrating models of recovery in addiction and mental health, <i>J of Dual Diagnosis</i> , 4 (3), 2008, 273-290. | USA | Literature review and consultation (n=45 people with addictions or in recovery from serious mental illness. | 8/25 |
| 52 | Sullivan W. A long and winding road: The process of recovery from severe mental illness, in Spaniol L, Gagne C, Koehler M, (eds) <i>Psychological and social aspects of Psychiatr disabilities</i> , Boston University Center, 1997. | USA | Semi-structured interviews (n=46 current and former service users) | 13/25 |
| 53 | Mancini M. Consumer-providers' theories about recovery from serious psychiatric disabilities, chapter 2, from Rosenberg, <i>Community Mental Health: Challenges for the 21st Century</i> , Routledge, 2006. | USA | Semi-structured interviews (n==15 people diagnosed with a psychiatric disability who also provide peer-support services to others) | 11/25 |
| 54 | Ridge D, Ziebland S. "The old me could never have done that": how people give meaning to recovery following depression, <i>Qual. Health Research</i> , 2006, 16 (8), 1038-1053. | UK | Open-ended interviews (n=38 people who have had depression) | CHECK RATS |
| 55 | Sydney West Area Health Service, (2008) Maintaining wellness and promoting recovery, sections 4-6, in <i>The wellness guide – a resource to support the recovery journey</i> , March 2008. | Australia | Part of a Wellness Guide developed in partnership between consumers and clinicians. | |
| 56 | Armstrong N, Steffen J. The Recovery Promotion Fidelity Scale: Assessing the organizational promotion of recovery, <i>Community Ment. Health J</i> , 2009, 45 (3), 163-170. | USA | Literature review and concept mapping (n= 5 focus groups) and survey | 16/25 |
| 57 | Noordsy D, Toeey W, Mueser K, O'Keefe C, Fox L. Recovery from severe mental illness: an intrapersonal and functional outcome definition, <i>Int. Review of Psychiatry</i> , 2002, 14 , 318-326. | USA | Focus groups and observation | Not rated |
| 58 | Forchuk C, Jewell J, Tweedell D, Steinnage IL. Reconnecting the client experience of recovery from psychosis, <i>Perspectives in Psychiatr Care</i> , 2003, 39 (4) 141-150. | Canada | Interviews and observation (n=10 patients over the initial year of treatment with clozapine or risperidone) | 16/25 |
| 59 | Baxter E, Diehl S. Emotional stages: Consumers and family members recovering from the trauma of mental illness, <i>Psychiatr Rehab. J</i> , 1998, 21 (4), 349-355. | USA | Interviews (n=40 consumers) | 11/25 |
| 60 | Oades L, Deane F, Crowe T, Lambert W, Kavanagh D, Lloyd C. Collaborative recovery: An integrative model for working with individuals who experience chronic and recurring mental illness. <i>Australasian Psychiatry</i> , 2005, 13 (3), 279-284. | Australia | Multi-site study in 9 organisations | Not rated |
| 61 | Glover H. Lived experience perspectives, in <i>Handbook of psychosocial rehabilitation</i> , King R, Lloyd C, Meehan T, Wiley-Blackwell, 2007. | Australia | Literature review and personal narrative | Not rated |
| 61 | Ridgeway P. Re-Storying psychiatric disability: Learning from first person recovery narratives, <i>Psychiatr Rehab. J</i> , 2001, 24 (4), 335-343 | USA | Grounded theory analysis of 4 existing 'seminal' narratives | 17/25 |
| 63 | Bonney S, Stickley T. Recovery and mental health: A review of the British literature, <i>J of Psychiatr and Ment. Health Nurs.</i> , 2008, 15 (2), 140-153. | UK | Literature review | Not rated |
| 64 | Mead S, Copeland M. What recovery means to us: Consumers' perspectives, | USA | Personal narratives and opinion piece | 8/25 |

| | | | | |
|----|---|-------------|--|-----------|
| | <i>Community Ment. Health J</i> , 2000, 36 (3), 315-328. | | | |
| 65 | Sowers W. Transforming systems of care: the American Association of Community Psychiatrists guidelines for recovery oriented services, <i>Community Ment. Health J</i> , 2005, 41 (6), 757-774 | USA | Literature review | Not rated |
| 66 | Plum K. How patients view recovery: what helps, what hinders, <i>Archives of Psychiatr Nurs.</i> , 1987, 1 (4), 285-293. | USA | Analysis of narratives (n=20) | 13/25 |
| 67 | Ahern L, Fisher D. Recovery at your own PACE (Personal Assistance in Community existence). <i>J of Psychosocial Nurs. & Ment. Health Services</i> , 2001, 39 (4), 22-32. | USA | Literature review and qualitative research | Not rated |
| 68 | Jacobson N, Curtis L. Recovery as policy in mental health services: Strategies emerging from the states, <i>Psychiatr Rehab. J</i> , 2000, 23 (4), 333-341. | USA | Literature review | Not rated |
| 69 | Lunt A. A theory of recovery. <i>J of Psychosocial Nurs. & Ment. Health Services</i> , 2002, 40 (12), 32-39. | USA | Literature review and opinion piece | Not rated |
| 70 | Nicholls V. <i>Feeding the flowers: SPN perspective on recovery</i> , 2007. | UK | Literature review and qualitative research | Not rated |
| 71 | Ralph R, Risman J, Kidder, K. <i>The Maine contingent of the recovery advisory group</i> , May, 1999. | USA | Personal narratives and literature review | 6/25 |
| 72 | Mental Health Providers Forum, <i>The recovery star model</i> , 2008. | UK | Measure development | Not rated |
| 73 | Mental health providers forum, <i>The recovery star model and cultural competency</i> , BAME Pilot Report, 2009. | UK | Pilot study to test measure with BME population | Not rated |
| 74 | Brown M, Essien P, Etim-Ubah P et al. <i>Report of the community led research project focussing on male African and African Caribbean perspectives on recovery</i> , Southside Partnership Fanon, 2008. | UK | Semi-structured interviews and questionnaires (n=91) | 20/25 |
| 75 | Mancini M, Hardiman E, Lawson H. Making Sense of It All: Consumer Providers' Theories about Factors Facilitating and Impeding Recovery from Psychiatric Disabilities, <i>Psychiatr Rehab. J</i> , 2005, 29 (1), 48-55. | USA | Semi-structured interviews (n=15 adults recovering from serious psychiatric disability and leading consumer provision of mental health services) | 14/25 |
| 76 | Bradstreet S, Brown W. Elements of recovery: Int. learning and the Scottish context, <i>SRN Discussion Paper Series Report No. 1</i> . 2004. | UK | Literature review | Not rated |
| 77 | Jacobson N, Greenley D. (2001) What is recovery? A conceptual model and explication. <i>Psychiatr Services</i> , 52 (4), 482-485. | USA | Synthesis of consumer narratives | Not rated |
| 78 | Lapsley H, Waimarie Nikora L, Black R. <i>Kia Mauri Tau! Narratives of recovery from disabling mental health problems</i> . Report of the University of Waikato Mental Health Narratives Project. Wellington: Mental Health Commission, 2002. | New Zealand | Interviews (n=40 who once had a disabling mental health problem) | 20/25 |
| 79 | Jenkins J, Carpenter-Song E. The new paradigm of recovery from schizophrenia: cultural conundrums of improvement without cure, <i>Culture, Medicine and Psychiatry</i> , 2006, 29 (4), 379-414. | USA | Interviews (n=90 people attending community out-patient clinics) | 18/25 |
| 80 | Barton R. The rehabilitation-recovery paradigm: A statement of philosophy for a public mental health system, <i>Psychiatr Rehab. Skills</i> , 1998, 2 (2), 171-187. | USA | Literature review | Not rated |
| 81 | Spaniol L, Gagne C, Koehler M. The recovery framework in rehabilitation and mental health, chapter 4, in Moxley, D. & Finch, J. <i>Sourcebook of rehabilitation and mental health practice</i> . New York, US: Kluwer Academic/Plenum, 2003. | USA | Literature review | Not rated |
| 82 | Glover H. Recovery based service delivery: Are we ready to transform the words into a paradigm shift? <i>Australian e-J for the Advancement of Ment. Health</i> , 2005, 4 (3), 1-4. | Australia | Opinion piece | Not rated |

| | | | | |
|----|--|----------|--|-----------|
| 83 | Irish Mental Health Commission. <i>A recovery approach within the Irish mental health services: A framework for development</i> , 2008. | Ireland | Literature review | Not rated |
| 84 | Ochocka J, Nelson G, Janzen R. Moving Forward: Negotiating Self and External Circumstances in Recovery, <i>Psychiatr Rehab. J</i> , 2005, 28 (4), 315-322. | Canada | In-depth interviews (n=28 people who had experienced serious mental health problems) | 14/25 |
| 85 | Brown W. The possibility of wellness, <i>Ment. Health Today</i> , 2007, Sept. 23-26. | Scotland | Semi-structured interviews (n=64) | 12/25 |
| 86 | Steen M. Essential structure and meaning of recovery from clinical depression for middle-adult women: a phenomenological study, <i>Issues in Ment. Health Nurs.</i> , 1996, 17 (2), 73-92. | USA | Interviews (n=22 participants with clinical, unipolar depression) | 13/25 |
| 87 | Fisher D. Healthcare reform based on an empowerment model of recovery by people with psychiatric disabilities, <i>Hospital and community psychiatry</i> , 1994, 45 (9), 913-915. | USA | Opinion paper | Not rated |
| 88 | Substance Abuse and Mental Health Service Administration, <i>National Consensus statement on mental health recovery</i> , 2004. | USA | Consensus methods (n= 110 expert panellists) | Not rated |
| 89 | Onken S, Craig C, Ridgway P, Ralph R, Cook J. An analysis of the definitions and elements of recovery: A review of the literature. <i>Psychiatr Rehab. J</i> , 31 (1), 9-22. 2007 | USA | Literature review | Not rated |
| 90 | Anthony W. Recovery from mental illness: The guiding vision of the mental health service system in the 1990s, <i>Psychosocial Rehab. J</i> , 1993, 16 (4), 11-23. | USA | Opinion piece and literature review | Not rated |
| 91 | Pitt L, Kilbride M, Nothard S, Welford M, Morrison A. Researching recovery from psychosis: a user-led project, <i>Psychiatr Bulletin</i> , 2007, 31 , 55 - 60. | UK | User-led interview study (n= 7 people in recovery) | 19/25 |
| 92 | Anderson B, Munchel W. <i>Opportunity on the doorstep: recovery-oriented leadership</i> , 2005 Village ISA and Community Activators, Inc. | USA | Opinion piece | Not rated |
| 93 | Borg M, Davidson, L. The nature of recovery as lived in everyday experience, <i>J of Ment. Health</i> , 2008, 17 (2), 129-140. | Norway | Interviews (n=13 individuals in recovery) | 14/25 |
| 94 | Asmundsdottir E. Creation of New Services: Collaboration Between Mental Health Consumers and Occupational Therapists, <i>Occupational Therapy in Ment. Health</i> , 2009, 25 (2), 115-126. | Iceland | Interviews and focus groups (n=25) | 14/25 |
| 95 | Davis E, Velleman R, Smith G, Drage M. Psychosocial developments: Towards a model of recovery, in Velleman R, Davies E, Smith G, & Drage M. (eds.) <i>Changing outcomes in Psychosis, Collaborative cases from practitioners, users and carers</i> , pp1-21., BPS Blackwell, 2006. | UK | Literature review | Not rated |
| 96 | Warren K. <i>Exploring the concept of recovery from the perspective of people with mental health problems</i> . Norwich School of Social Work and Psychosocial Studies, University of East Anglia, 2003. | UK | Literature review, narrative analysis, interviews | 9/25 |
| 97 | Piat M, Sabetti J, Bloom D. The importance of medication in consumer definitions of recovery from serious mental illness: A qualitative study, <i>Issues in Ment. health Nurs.</i> , 2009, 30 (8), 482-490. | Canada | Semi-structured interviews (n= 54 consumers of mental health services) | 18/25 |

For full list of excluded studies, please contact Dr Mary Leamy (mary.leafy@kcl.ac.uk).

Table DS2 Vote counting of recovery processes

| Recovery processes | Number (%) of 87 studies |
|---|---------------------------------|
| Category 1: Connectedness | 75 (86%) |
| <i>1.1 Peer support and support groups</i> | 39 (45%) |
| 1.1.1 Availability of peer support | 22 (25%) |
| 1.1.2 Becoming a peer support worker or advocate | 17 (20%) |
| <i>1.2 Relationships</i> | 33 (38%) |
| 1.2.1 Building upon existing relationships | 19 (22%) |
| 1.2.2 Intimate relationships | 9 (10%) |
| 1.2.3 Establishing new relationships | 8 (9%) |
| <i>1.3 Support from others</i> | 53(61%) |
| 1.3.1 Support from professionals | 42 (48%) |
| 1.3.2 Supportive people enabling the journey | 27 (31%) |
| 1.3.3 Family support | 26 (30%) |
| 1.3.4 Friends and peer support | 18 (21%) |
| 1.3.5 Active or practical support | 4 (5%) |
| <i>1.4 Being part of the community</i> | 35 (40%) |
| 1.4.1 Contributing and giving back to the community | 21 (24%) |
| 1.4.2 Membership of community organisations | 13 (15%) |
| 1.4.3 Becoming an active citizen | 11 (13%) |
| Category 2: Hope and optimism about the future | 69 (79%) |
| <i>2.1 Belief in possibility of recovery</i> | 30 (34%) |
| <i>2.2 Motivation to change</i> | 15 (17%) |
| <i>2.3 Hope-inspiring relationships</i> | 12 (14%) |
| 2.3.1 Role-models | 8 (9%) |
| <i>2.3 Positive thinking and valuing success</i> | 10 (11%) |
| <i>2.4 Having dreams and aspirations</i> | 7 (8%) |
| Category 3: Identity | 65 (75%) |
| <i>3.1 Dimensions of identity</i> | 8 (9%) |
| 3.1.1 Culturally specific factors | 7 (8%) |
| 3.1.2 Sexual identity | 2 (2%) |
| 3.1.3 Ethnic identity | 4 (5%) |
| 3.1.4 Collectivist notions of identity | 6 (7%) |
| <i>3.2 Rebuilding/redefining positive sense of self</i> | 57 (66%) |
| 3.2.1 Self-esteem | 21 (24%) |
| 3.2.2 Acceptance | 21 (24%) |

| | |
|---|-----------------|
| 3.2.3 Self-confidence and self-belief | 11 (13%) |
| 3.3 <i>Over-coming stigma</i> | 40 (46%) |
| 3.3.1 Self-stigma | 27 (31%) |
| 3.3.2 Stigma at a societal level | 32 (37%) |
| <hr/> | |
| Category 4: Meaning in life | 59 (66%) |
| 4.1 <i>Meaning of mental illness experiences</i> | 30 (34%) |
| 4.1.1 Accepting or normalising the illness | 22 (25%) |
| 4.2 <i>Spirituality (including development of spirituality)</i> | 36 (41%) |
| 4.3 <i>Quality of life</i> | 57 (65%) |
| 4.3.1 Well-being | 27 (31%) |
| 4.3.2 Meeting basic needs | 18 (21%) |
| 4.3.3 Paid voluntary work or work related activities | 19 (22%) |
| 4.3.4 Recreational and leisure activities | 8 (9%) |
| 4.3.5 Education | 7 (8%) |
| 4.4 <i>Meaningful social and life goals</i> | 15 (17%) |
| 4.4.1 Active pursuit of previous or new life or social goals | 15 (17%) |
| 4.4.2 Identification of previous of new life or social goals | 8 (9%) |
| 4.5 <i>Meaningful life and social roles</i> | 40 (46%) |
| 4.5.1 Active pursuit of previous or new life or social roles | 40 (46%) |
| 4.5.2 Identification of previous of new life or social roles | 34 (39%) |
| 4.6 <i>Rebuilding of life</i> | 20 (23%) |
| 4.6.1 Resuming with daily activities and daily routine | 12 (14%) |
| 4.6.2 Developing new skills | 8 (9%) |
| <hr/> | |
| Category 5: Empowerment | 79 (91%) |
| 5.1 <i>Personal responsibility</i> | 79 (91%) |
| 5.1.1 Self-management | 60 (69%) |
| Coping skills | 25 (29%) |
| Managing symptoms | 22 (25%) |
| Self-help | 12 (14%) |
| Resilience | 25 (29%) |
| Maintaining good physical health and well-being | 12 (14%) |
| 5.1.2 Positive risk-taking | 17 (20%) |
| 5.2 <i>Control over life</i> | 78 (90%) |
| 5.2.1 Choice | 31 (36%) |
| Knowledge about illness | 17 (20%) |
| Knowledge about treatments | 7 (8%) |
| 5.2.2 Regaining independence and autonomy | 23 (26%) |

| | |
|--|--------------|
| 5.2.3 Involvement in decision-making | 23(26%) |
| Care planning | 35 (40%) |
| Crisis planning | 7 (8%) |
| Goal setting | 12 (14%) |
| Strategies for medication | 25 (29%) |
| Medication not whole solution | 11 (13%) |
| 5.2.4 Access to services and interventions | 13 (15%) |
| 5.3 <i>Focussing upon strengths</i> | 14 (16%) |

BJPpsych

The British Journal of Psychiatry

Conceptual framework for personal recovery in mental health: systematic review and narrative synthesis

Mary Leamy, Victoria Bird, Clair Le Boutillier, Julie Williams and Mike Slade
BJP 2011, 199:445-452.

Access the most recent version at DOI: [10.1192/bjp.bp.110.083733](https://doi.org/10.1192/bjp.bp.110.083733)

**Supplementary
Material**

Supplementary material can be found at:
<http://bjp.rcpsych.org/content/suppl/2013/11/21/199.6.445.DC1>

References

This article cites 25 articles, 5 of which you can access for free at:
<http://bjp.rcpsych.org/content/199/6/445#BIBL>

**Reprints/
permissions**

To obtain reprints or permission to reproduce material from this paper, please
write to permissions@rcpsych.ac.uk

**You can respond
to this article at**

[/letters/submit/bjprcpsych;199/6/445](http://bjp.rcpsych.org/letters/submit/bjprcpsych;199/6/445)

**Downloaded
from**

<http://bjp.rcpsych.org/> on August 17, 2017
Published by [The Royal College of Psychiatrists](http://www.rcpsych.ac.uk)
