

Choice and rejection of psychiatry as a career: surveys of UK medical graduates from 1974 to 2009

Michael J. Goldacre, Seena Fazel, Fay Smith and Trevor Lambert

Background

Recruitment of adequate numbers of doctors to psychiatry is difficult.

Aims

To report on career choice for psychiatry, comparing intending psychiatrists with doctors who chose other clinical careers.

Method

Questionnaire studies of all newly qualified doctors from all UK medical schools in 12 qualification years between 1974 and 2009 (33 974 respondent doctors).

Results

One, three and five years after graduation, 4–5% of doctors specified psychiatry as their first choice of future career. This was largely unchanged across the 35 years. Comparing intending psychiatrists with doctors who chose other careers, factors with a greater influence on psychiatrists' choice included their experience of the subject at medical school, self-appraisal of their own skills, and inclinations before medical school. In a substudy of doctors who initially

considered but then did not pursue specialty choices, 72% of those who did not pursue psychiatry gave 'job content' as their reason compared with 33% of doctors who considered but did not pursue other specialties. Historically, more women than men have chosen psychiatry, but the gap has closed over the past decade.

Conclusions

Junior doctors' views about psychiatry as a possible career range from high levels of enthusiasm to antipathy, and are more polarised than views about other specialties. Shortening of working hours and improvements to working practices in other hospital-based specialties in the UK may have reduced the relative attractiveness of psychiatry to women doctors. The extent to which views of newly qualified doctors about psychiatry can be modified by medical school education, and by greater exposure to psychiatry during student and early postgraduate years, needs investigation.

Declaration of interest

None.

Many countries have difficulty in recruiting adequate numbers of doctors to psychiatry. In the USA, it has been estimated that three-quarters of its counties (the administrative level within US states) have a shortage of psychiatric prescribers.¹ The percentage of medical students pursuing a psychiatry residency in the USA was 7–10% in the 1940s but it dropped to 3–4% in 2002–2007.^{2,3} In England, in 2009, 15% of all unfilled consultant posts were in psychiatry;⁴ and a census by the Royal College of Psychiatrists concluded that recruitment to specialty training posts remains difficult.⁵ The UK Medical Careers Research Group (UKMCRG) has studied the career preferences of junior doctors and their subsequent career pathways for many years.⁶ Here, we update our findings on career choice for psychiatry and report on reasons given by doctors for choosing psychiatry. In addition, unlike our previous study, we have added new findings examining doctors who told us that they had initially considered a career in psychiatry and then decided not to pursue it.

Method

The UKMCRG has undertaken cohort studies of all medical graduates who qualified in the years 1974, 1977, 1980, 1983, 1993, 1996, 1999, 2000, 2002, 2005, 2008 and 2009 from all UK medical schools. Our methods have been described in detail elsewhere.^{7,8} Near the end of the first, third and fifth years after qualification (and at longer time intervals after that), postal questionnaires are sent to all the graduates. The starting point for each survey is the cohort as it originally was at the time of

qualification. Current addresses are supplied to us by the UK General Medical Council and by the doctors themselves from replies to successive surveys. In calculating our response rate, we excluded those known to have died, those who expressed a wish not to participate and those whose address was untraceable. Others in the original cohort who did not initially reply were sent several reminders.

We ask graduates 'What is your choice of long-term career?' This is an open response question, inviting the doctors to name their preferred clinical specialty or any other choice of career. We invite them to list up to three specialty choices. If they specify more than one specialty, they are asked to indicate whether their choices are of equal priority (termed 'tied' choices by us). In this paper, we counted first choices for psychiatry tied with another specialty (as well as untied first choices) as doctors choosing psychiatry. We also ask them to indicate whether their choice of specialty is 'definite', 'probable' or 'uncertain'.

The graduates of 1993, 1996, 1999, 2000, 2002, 2008 and 2009 were asked to indicate whether each factor from a list of 11 had influenced their choice of specialty 'not at all', 'a little' or 'a great deal'. In addition to comments about reasons for choosing their specialty, some doctors provided spontaneous comments to us about career choices they had considered and rejected. Accordingly, we decided to add a structured question on this in the survey of the doctors who qualified in 2002, 2005 and 2008: 'Is there a choice of long-term career which you have seriously considered but have now decided not to pursue?' If they answered yes, they were asked 'What was that choice?' and 'What are your most important reasons for rejecting that choice?' Reasons for

rejection were sought as free-text responses. Some respondents gave a single reason for rejection; others gave several reasons. We read the responses and developed a coding scheme to reflect the main themes raised by respondents (Appendix 1). Two coders then independently coded all the reasons given, by theme, and compared each other's coding. Any differences between coders were resolved through discussion. We compared findings from doctors who specified psychiatry with those from doctors who reported that they had considered but not pursued other clinical specialties.

We used χ^2 statistics (reporting Yates's continuity correction where there was only one degree of freedom, and Mantel-Haenszel linear-by-linear χ^2 -test for linear association between two variables) correlation analysis to make comparisons between groups of doctors and to test their statistical significance. A binary logistic regression model was used to investigate variation between medical schools in choice for psychiatry: categorical predictors were gender, cohort and medical school, and the dependent variable was whether or not psychiatry had been chosen as a first, second or third choice.

This study was approved by the UK National Research Ethics Service (ref 04/Q1907/48).

Results

The survey questionnaires were sent to 51 538 doctors covering all 12 cohorts in year 1; and 33 974 (65.9%) replied. Three years after qualification, the survey questionnaire was sent to 38 131 doctors covering the first ten cohorts: 26 174 (68.6%) replied. Five years after qualification, the survey questionnaire was sent to 29 111 doctors covering the first eight cohorts (1974–1980 and 1993–2002: the 1983 cohort was not surveyed in year 5) and 20 239 (69.5%) replied.

Early career choices for psychiatry: first choices

Considering all cohorts combined, 1 year after qualification 4.2% of all respondents (4.1% of men and 4.3% of women) indicated that their first choice of long-term career was psychiatry

($\chi^2_1=0.8$, $P>0.05$, comparing men and women; Table 1, online Fig. DS1 and online Table DS1). There was no overall linear trend over time (across cohorts) in career choice for psychiatry (Table 1, Fig. DS1 and Table DS1).

Three years after qualification, 4.7% of all respondents (4.4% of men and 5.1% of women) gave psychiatry as their first choice, as did 5.0% of all respondents 5 years after (4.5% of men and 5.4% of women). Three and five years after qualification, the percentage of women who chose psychiatry was significantly higher than that of men ($\chi^2_1=8.7$, $P<0.01$, comparing men and women, in each year). Although there was no significant linear relationship in career choice across cohorts overall, from about 1999 the percentage of women who chose psychiatry in years 3 and 5 started to decline (χ^2_1 for linear trend 14.2, $P<0.001$ and $\chi^2_1=10.2$, $P<0.01$ respectively; Table 1, Fig. DS1 and Table DS1).

Early career choices for psychiatry: first, second and third choices

Considering first, second and third choices of career combined, 7.5% of all respondents (7.3% of men and 7.6% of women) indicated that they were at least considering a choice for psychiatry in year 1 ($\chi^2_1=0.9$, $P>0.05$, comparing men and women; Table 1, Fig. DS1 and Table DS1). Three years after graduation, 6.8% of respondents (6.1% of men, 7.5% of women) gave psychiatry as a possible choice ($\chi^2_1=19.3$, $P<0.001$, comparing men and women). Five years after, 6.1% of respondents (5.6% of men, 6.6% of women) gave psychiatry as a choice ($\chi^2_1=8.3$, $P<0.01$, comparing men and women).

Considering all choices for psychiatry – first, second and third choices combined – there was a significant downward linear trend over time ($P<0.001$ for choices given at years 1 and 3, $P<0.01$ for choices at year 5). There was a steeper decline in women than men. Psychiatry had been a more popular choice among women than men in the earlier cohorts; and the decline from early to late cohorts was greater in women than men (Table 1, Fig. DS1 and Table DS1). Against the general trend in first-year choices, there was a slight upturn in the 2008 and 2009 cohorts in year 1 (Table 1, Fig. DS1 and Table DS1).

Table 1 Percentages of respondents choosing psychiatry as their first, second or third choice of career at 1 (1974–2009), 3 (1974–2005) and 5 (1974–2000) years after graduation^a

Cohort	Percentages choosing psychiatry as first choice									Percentages choosing psychiatry as first, second or third choice								
	Year 1			Year 3			Year 5			Year 1			Year 3			Year 5		
	Men	Women	Total	Men	Women	Total	Men	Women	Total	Men	Women	Total	Men	Women	Total	Men	Women	Total
1974	3.8	4.9	4.1	3.5	5.1	4.0	3.8	5.8	4.4	8.7	11.7	9.5	5.4	8.2	6.2	4.8	8.9	5.9
1977	3.6	6.0	4.4	3.4	7.4	4.8	3.7	7.3	4.9	7.3	10.8	8.5	5.5	11.4	7.5	5.8	9.5	7.1
1980	3.5	3.9	3.7	5.2	5.5	5.3	4.9	6.5	5.5	7.4	8.2	7.7	7.2	8.4	7.6	6.5	8.4	7.2
1983	4.2	5.1	4.6	4.3	6.6	5.3	n/a ^b	n/a ^b	n/a ^b	8.8	12.5	10.3	6.8	10.7	8.3	n/a ^b	n/a ^b	n/a ^b
1993	4.6	4.3	4.4	3.8	5.1	4.5	4.3	5.7	5.0	7.4	7.8	7.6	5.4	7.6	6.5	4.9	6.9	5.9
1996	4.2	4.2	4.2	5.1	4.6	4.9	5.2	4.7	5.0	7.2	6.7	7.0	6.8	6.2	6.5	6.1	5.6	5.8
1999	4.6	4.9	4.7	4.5	5.5	5.1	4.8	5.9	5.4	6.7	7.3	7.1	5.3	8.0	6.8	5.6	6.4	6.1
2000	4.4	3.8	4.1	5.3	5.1	5.2	5.6	4.8	5.1	6.3	5.8	6.0	7.0	7.3	7.2	6.2	5.7	5.9
2002	4.6	3.9	4.2	3.6	4.5	4.1	4.2	4.1	4.2	7.1	6.1	6.5	4.4	5.8	5.3	4.3	4.5	4.4
2005	3.7	3.3	3.5	4.5	3.6	3.9	n/a ^b	n/a ^b	n/a ^b	6.2	5.7	5.9	6.2	4.8	5.4	n/a ^b	n/a ^b	n/a ^b
2008	4.9	4.0	4.3	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	7.7	6.6	7.0	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b
2009	3.3	4.6	4.2	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	5.7	8.3	7.4	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b
Total	4.1	4.3	4.2	4.4	5.1	4.7	4.5	5.4	5.0	7.3	7.6	7.5	6.1	7.5	6.8	5.6	6.6	6.1

a. Numbers of respondents were: 16 488 (men), 17 486 (women) and 33 974 (total) in year 1; 13 429 (men), 12 745 (women) and 26 174 (total) in year 3; and 10 544 (men), 9 695 (women) and 20 239 (total) in year 5.

b. Third- and fifth-year choices not yet available for graduates of 2008/9; fifth-year choices not yet available for graduates of 2005; the 1983 cohort was not surveyed in year 5.

Definite, probable and uncertain choices

We compared certainty of choice between aspiring psychiatrists and those who gave their first choice as general practice or a non-psychiatric hospital specialty (Table 2 and online Table DS2). In years 1, 3 and 5, those who chose psychiatry were less certain about their career preference than those who chose general practice, but more certain than doctors who chose other hospital specialties.

Choices by medical school

Choices for psychiatry varied little by medical school attended (online Table DS3). Comparing year-1 choices for the 23 schools on which we had more than 1 year's data, and taking a cut-off of $P=0.01$ for significance, one school was significantly high and two were significantly low (Table DS3). At year 3, no school was significantly high or low. At year 5, one school was significantly low and two were significantly high. There was an indication in the results that a slightly higher percentage of graduates from Scotland and Northern Ireland favoured a career in psychiatry (Table DS3).

Factors influencing career choice

In year 1, three-quarters of respondents rated 'enthusiasm/commitment: what I really want to do' as influencing their choice of psychiatry a great deal (Table 3 and online Table DS4). 'Self-appraisal of own skills', 'student experience of subject' and

'hours/working conditions' were rated as having a great deal of influence by over half of respondents. 'Self-appraisal of own skills', 'student experience of subject' and 'inclinations before medical school' were rated as having a 'great deal of influence' by a higher percentage of aspiring psychiatrists than of those who chose general practice or other hospital specialties. 'Hours/working conditions' were rated as influential by a higher percentage of doctors who chose psychiatry than those who chose other hospital specialties. Women who chose psychiatry rated 'domestic circumstances' as having influenced their choice a 'great deal' more than did men (33% of women, 18% of men; $\chi^2_1=20.3$, $P<0.001$). Women who chose psychiatry rated 'hours/working conditions' as more influential on their career choice than did men (62% of women, 54% of men; $\chi^2_1=5.1$, $P<0.05$). Men who chose psychiatry rated 'promotion/career prospects' as influential, more so than did women (33% of men, 25% of women; $\chi^2_1=5.1$, $P<0.05$).

By the third year after graduation, 'experience of jobs so far' was rated as the most influential factor. Women who chose psychiatry rated this as having had more influence than did men (77% of women, 63% of men; $\chi^2_1=6.9$, $P<0.01$). By the fifth year 'enthusiasm/commitment' was rated most important. Women who chose psychiatry continued to rate 'experience of jobs so far' as influential in higher numbers than did men (73% of women, 65% of men; $\chi^2_1=4.2$, $P<0.05$). 'Student experience of subject' was more important for those choosing psychiatry than those who chose general practice or other hospital specialties across all three periods after qualification.

Table 2 Comparison between doctors who chose psychiatry, general practice or other clinical specialties: percentages of doctors who specified that they were 'definite', 'probable' or 'uncertain' about their first choice of long-term career, at 1, 3 and 5 years after graduation

Firmness of choice	Percentages								
	Year 1			Year 3			Year 5		
	Psychiatry	General practice	Other	Psychiatry	General practice	Other	Psychiatry	General practice	Other
Definite	30.0	36.1	25.9	54.1	58.8	42.0	65.5	72.9	67.3
Probable	48.0	44.6	51.4	37.3	34.0	44.8	29.6	23.3	28.1
Uncertain	22.0	19.3	22.7	8.6	7.2	13.1	4.9	3.8	4.6
Total	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table 3 Percentages of doctors who specified each factor as influencing their choice of long-term career a great deal: graduates of 1993, 1996, 1999, 2000, 2002, 2008 and 2009^a

Factor	Year 1			Year 3			Year 5		
	Psychiatry (n = 869)	General practice (n = 5615)	Other ^b (n = 12 829)	Psychiatry (n = 655)	General practice (n = 4020)	Other ^b (n = 8265)	Psychiatry (n = 651)	General practice (n = 4422)	Other ^b (n = 7448)
Domestic circumstances	26.4 ^c	49.0***	15.5***	24.1	48.7***	15.3***	36.5	68.5***	26.9***
Hours/working conditions	58.5 ^c	75.1***	28.6***	58.1	81.5***	33.9***	64.6	87.7***	39.4***
Eventual financial prospects	9.4	14.9***	11.9	9.2	22.8***	11.2	6.0 ^c	17.7***	9.7
Promotion/career prospects	28.9 ^c	16.2***	23.9	28.0	22.8	24.5	27.0	20.6***	25.1
Self-appraisal of own skills	63.8	47.7***	49.5***	67.1	49.8***	50.0***	71.5	52.6***	58.3***
Advice from others	10.7	15.5	17.0***	12.4	16.6	17.6	9.5	11.5	15.4***
Student experience of subject	58.7	38.3***	48.1***	36.7	20.8***	27.0***	36.1	19.8***	24.4***
Particular teacher/department	23.3	13.3***	32.7***	15.9	11.5	33.3***	14.4	6.0***	30.4***
Inclinations before medical school	22.4	15.1***	13.7***	18.3	14.8	12.1	17.0	13.0	9.5***
Experience of jobs so far	37.5	47.1***	54.5***	71.3 ^c	54.0***	69.7	69.2 ^c	53.2***	72.2
Enthusiasm/commitment: what I really want to do	75.7	63.5***	73.0	67.4	56.5***	67.2	78.9	63.2***	80.5

a. The numbers for this table are provided in online Table DS4.

b. Other hospital specialties (i.e. all hospital specialties other than psychiatry).

c. Significant difference between men and women within the group that chose psychiatry.

*** $P<0.001$ when compared with psychiatry.

Doctors who considered psychiatry as a career and then decided not to pursue it

This part of the study was based on questionnaires that had been sent to 16 361 doctors who had qualified in 2002, 2005 or 2008. The percentage of respondents who specified that they had seriously considered but then not pursued a specialty were very similar in the successive cohorts: this was specified by 29% (797/2778) of graduates in 2002, 27% (843/3128) in 2005, and 29% (933/3249) in 2008. After exclusions of statements that could not be coded, and doctors whose switches of specialty were within the same broader specialty group (e.g. between different branches of hospital medicine or surgery, as described elsewhere⁹), there were 2267 respondents – 811 men and 1456 women – who had considered and rejected a specialty and who gave at least one classifiable reason for doing so (25% of the total number of respondents).

Of the 2267 respondents, 99 (4.4%) specified that they had considered and then decided not to pursue psychiatry (4.2% of men, 4.5% of women, a very similar percentage to those who chose it, Table 1). There was no significant upward or downward trend across the cohorts in the percentage that considered and rejected psychiatry: it was 4.8% in the 2002 cohort, 3.7% in 2005 and 4.7% in 2008.

Reasons for rejecting psychiatry, after having seriously considered it as a career

We compared the reasons for rejecting a specialty given by doctors who rejected psychiatry with those given by the 2168 doctors who had seriously considered but then decided not to pursue other specialties. Statements that we classified as ‘job content’ were given by 71% of the doctors who rejected psychiatry and by 33% of doctors who rejected other specialties (Table 4 and online Table DS5).

Illustrative examples of comments that we coded as ‘job content’ are reproduced, exactly as written by the doctors, in Appendix 2.

Twelve per cent of doctors who rejected psychiatry and four per cent of those who rejected other specialties gave reasons that we classified as ‘self-appraisal’ (Appendix 2). All other reasons, such as those relating to competition, training and working relationships within the specialty were mentioned by less than

10% of doctors rejecting psychiatry. ‘Work–life balance’ was the most frequently mentioned reason given by those rejecting specialties other than psychiatry (41%), but was only mentioned by 5% of those rejecting psychiatry.

Choosing psychiatry, following rejection of another specialty

Of the doctors who had considered and rejected a specialty other than psychiatry, 4% then chose psychiatry. This is exactly in line with doctors’ general level of choice for psychiatry (Table 1).

Discussion

Early career choices for psychiatry

In the UK, until recently, newly qualified doctors undertook a ‘house officer’ year of working in a medical post and a surgical post under supervision, and then undertook a variable number of posts (depending on the wishes of individual doctors and their success in applications) before entering training for their choice of career specialty. A new programme of postgraduate medical training was introduced in the UK in 2005 which replaced the single ‘house officer’ year with two ‘foundation years’ before specialising. Prior to 2005, doctors could make their final career choice several years after graduation. It is now expected that most doctors will choose their broad area of specialty training, such as psychiatry, in their second postgraduate year so that they can start specialty training in their third year after graduation.

One year after graduation, about 4% of doctors chose psychiatry as their first choice of eventual career, similar to the 3–4% of doctors entering psychiatry in the USA between 1992 and 2002¹⁰ and 4% entering US psychiatry residency training in 2007.³ In the UK, the percentage of men who chose psychiatry has been consistent over the past 25 years. Previously, a higher percentage of women than men doctors in the UK have become psychiatrists. However, there has been a decline in choices for psychiatry made by women over the past decade, with a small increase in the last two surveys. Hours of work in medicine have been falling in the UK, following the implementation of European working time regulations, and working practices are considered to have become more ‘family friendly’ – aspects of working as a psychiatrist that have made it popular for women.¹¹ These

Table 4 Percentage distribution of reasons given for rejecting psychiatry and for rejecting all other specialties combined^a

Reason for rejection of specialty	Psychiatry (n=99), %	Other specialties (n=2168), %	Test ^b	P
Job content	71.7	33.3	$\chi^2_1=59.75$	<0.001
Self-appraisal	12.1	4.2	$\chi^2_1=12.2$	<0.001
Too stressful/lack of support/working conditions	8.1	10.3		
Work–life balance	5.1	41.9	$\chi^2_1=51.87$	<0.001
Training	2	8.3	$\chi^2_1=4.29$	<0.05
Training too long	2	5.3		
Future of the specialty uncertain	2	2		
Advice from others	2	1.4		
Working relationships in the specialty	2	5.2		
Lack of exposure and opportunities so far	2	2.1		
Competition	1	12.7	$\chi^2_1=12.7$	<0.001
Inadequate income	1	1.2		
Personal circumstances	1	1.5		
Fear of litigation	0	2.5		

a. Some doctors gave more than one reason and we counted each reason.
b. Significance tests, comparing those rejecting psychiatry and those rejecting other specialties.

changes to medicine will probably make some specialties more attractive to women doctors in the future than they have been in the past. If psychiatry loses its historical competitive edge with women, recruitment overall may worsen.

We have shown elsewhere that there is a high level of correspondence between early career choices for psychiatry and eventual destinations in the specialty.¹² In that publication, we showed that 75% who chose psychiatry in the first year after qualification eventually practised in it, as did 93% and 97% of those who specified it as their first choice at 3 and 5 years respectively after qualification. An early career choice for psychiatry is therefore highly predictive of an eventual career in the specialty.

Influences of medical school

A recent review of the impact of undergraduate experiences on recruitment into psychiatry highlighted an enthusiasm for psychiatry in school that dissipates once the student is in medical school.¹³ These authors listed factors which may dissuade medical students from choosing psychiatry. These included insufficient exposure to psychiatry in medical school, and a lack of status for psychiatry among peers. It has been suggested that efforts to increase the number of doctors choosing psychiatry might focus on medical schools which produce a smaller than average percentage of doctors wanting to pursue psychiatry.¹⁴ In our study, differences between medical schools were not particularly striking, although a higher percentage of doctors from medical schools in Scotland and Northern Ireland seem attracted to psychiatry. The extent to which these higher percentages reflect the characteristics of students selected at entry to medical school, and/or the influence of medical education, needs further examination. The degree to which different medical schools within other countries vary in their rates of recruitment into psychiatry is also an area for further study.

Factors influencing career choice for psychiatry

The factor cited as having a great deal of influence by the highest percentage of intending psychiatrists in year 1 was 'What I really want to do'. This was specified by 76% of intending psychiatrists, and it was a significantly higher level of enthusiasm for their specialty than that expressed by intending doctors for general practice. Three factors had a greater influence on career choice for psychiatry than on doctors' choice for other clinical careers: the doctors' self-appraisal of their own skills, their inclinations before medical school (which is not scored highly for most specialties), and their experience of the subject as a student. These factors in combination indicate, at least for some choosing psychiatry, an early and strong interest in the specialty.

Professional bodies have suggested offering more work experience placements to secondary school students (i.e. those aged 16–18) as a way to improve recruitment into psychiatry.¹⁴ Doctors' 'experience of the job so far' was less influential in year 1 for those choosing psychiatry than for those choosing other careers. The lack of work experience in the first year after medical graduation highlights the need for more doctors to spend time in a psychiatry placement in their early postgraduate years.¹⁵ Over the period covered by these studies, particularly in the early period, policies were implemented in the UK to emphasise community care for psychiatric patients and to reduce hospital in-patient admissions. This may have reduced the scope for junior doctors to gain experience of work in psychiatry in the hospital setting, and the change in work profile in psychiatry from in-patient to non-in-patient care may have had an impact on

junior doctors' predilection for the specialty in ways that we cannot assess.

A study by Dein *et al*¹⁶ asked consultant psychiatrists to specify the most important factors in their choice of career. The most important factors were empathy for patients, working conditions and experience of the subject as a student. However, their study included only psychiatrists and there was no element of comparison with other specialties. An Australian study of medical students found that the most important factors in career choice for psychiatry were that it was considered to be an interesting subject and intellectually challenging.¹⁷

Factors deterring career choice for psychiatry: those who once considered it

On the one hand, it is important that medical students and junior doctors are given adequate early experience of psychiatry. On the other hand, exposure may discourage some doctors. Positive choices for psychiatry, in our study, were heavily influenced by the nature of the specialty, the doctors' experience of it, and the doctors' self-appraisal of whether psychiatry is the right specialty for them. However, a similar range of factors – job content of the specialty, experience of it, and reflection on it as a career – were also cited as negative factors that influenced some doctors who seriously considered psychiatry as a career but then decided not to pursue it. It seems likely that views about psychiatry may be more polarised than those for other specialties.

Another UK study of doctors who initially chose psychiatry but then pursued a different career found that lack of respect among peers and the public, perceived threats of violence from patients, lack of resources and low morale were dissuading factors.¹⁸ A survey of Australian medical students found that the least attractive aspects of psychiatry included its lack of prestige and 'perceived absence of a scientific foundation'.¹⁷ These negative perceptions of psychiatry have been cited elsewhere^{19,20} and should be countered with real and good experiences of working in psychiatry during and after medical school. More importantly, the extent to which these factors can be altered by medical school education needs further investigation.

Future prospects

Although recruitment into psychiatry is difficult worldwide, demand for psychiatric services is increasing.²¹ The percentage of newly qualified doctors in the UK who choose psychiatry is below the level needed to replace the current generation of practising psychiatrists.^{4,22} The number of UK applicants for training in psychiatry is lower than that for any other large specialty,²³ and psychiatry in the UK is heavily reliant on the recruitment of international medical graduates. As in the UK, in the USA a substantial percentage of applications for psychiatry training/residency posts come from international medical graduates.³

One of our main findings is that the proportion of medical students choosing psychiatry as a career has not changed appreciably since 1974. This is despite a decade of initiatives by the Royal College of Psychiatrists to increase recruitment into psychiatry. These initiatives, mostly targeted at medical students, have principally aimed to emphasise the biopsychosocial model in psychiatry, and have highlighted how psychiatry is one of a number of professions contributing to the care of psychiatric patients.²⁰ In contrast, a group of mostly academic psychiatrists²⁴ recently called for more emphasis on the medical model in the teaching of psychiatry, and argued that this would contribute to improving the number and quality of candidates for psychiatry

training. Commentators have suggested that a lack of emphasis on medical models has fuelled perceptions that psychiatry may not have enough medical content to be attractive to many doctors.²⁵ Although these views were not widely shared among psychiatrists, the extent to which changes to teaching and training of students and junior doctors in psychiatry alters perceptions about it, and choices for it, needs testing. Our qualitative data provide some support for the perceived lack of applicability of the medical model as a reason for rejecting psychiatry after having seriously considered it as a career option.

We found that experiences of psychiatry as a student and in early medical training were important in determining choices. Student experience and experience of jobs so far seemed to be more influential for psychiatry than is the case for general practice and other specialties. This suggests that provision of more experience of psychiatry in foundation years might encourage recruitment to psychiatry. In keeping with the role of student experience, we also found potentially important differences in the proportion choosing psychiatry by medical school, but also differences in retention rates by medical school. These findings underscore the importance of student and early training experiences.

Strengths and limitations

The study has several strengths: it covers 12 cohorts of newly qualified doctors from all UK medical schools, going as far back as 1974; it has a high response rate for this type of study; it can be considered as nationally representative of the UK; and it is a prospective cohort study and therefore not subject to recall bias about early career intentions. Career intentions, as they were in years 1, 3 and 5, were provided by the doctors contemporaneously. Its longitudinal design also means that doctors' replies 1 year post-qualification can be compared with their later replies in years 3 and 5.

The study's limitations include the fact that it is restricted to UK medical graduates. There are 2 years post-graduation in which UK doctors work mostly in general medicine and surgery, and the length of this type of early training/internship differs by country. The importance of early choice is likely to be stronger in countries such as the USA where residency choice is made earlier. There are a few gaps in our data coverage: for example, the absence of year-1 data between 1983 and 1993 is because the funding body, the Department of Health, ceased funding the surveys in this period. Non-responder bias is possible, as with all surveys. We did not ask those who did not choose psychiatry whether certain factors might have made them choose it. We asked individuals about preferences, but some who choose psychiatry may not get posts in it.

Michael J. Goldacre, FFPH, FRCP, UK Medical Careers Research Group, Unit of Health-Care Epidemiology, Department of Public Health, University of Oxford; **Seena Fazel**, FRCPsych, **Fay Smith**, PhD, Department of Psychiatry, University of Oxford, Warneford Hospital, Oxford; **Trevor Lambert**, MSc, UK Medical Careers Research Group, Unit of Health-Care Epidemiology, Department of Public Health, University of Oxford, UK

Correspondence: Professor Michael J. Goldacre, UK Medical Careers Research Group, Department of Public Health, University of Oxford, Old Road Campus, Oxford OX3 7LF, UK. Email: michael.goldacre@dph.ox.ac.uk

First received 27 Feb 2012, final revision 4 Jul 2012, accepted 31 Jul 2012

Funding

S.F. is funded by the Wellcome Trust. This is an independent study commissioned and funded as part of a wider work programme by the Policy Research Programme in the Department of Health. The views expressed are not necessarily those of the funding body.

Appendix 1

Method for coding reasons for rejection of a specialty by theme

1. Each doctor's record includes a separate field for chosen specialty, rejected specialty (if any) and text comments on reasons for rejection, as well as for many other data items.
2. We selected records with a free-text comment about considering but then rejecting a specialty, for all specialties (not just psychiatry); and created a separate file containing the free text (and no other information). Accordingly, we were masked, in theme-scoring the comments, to the actual chosen specialty and the rejected specialty (although, of course, the doctors' text often mentioned them by name), gender, ethnicity and other personal data.
3. Two people independently read all the comments without a pre-conceived list of themes or coding schemes (i.e. using 'grounded' theory).
4. The themes were then agreed as those that had emerged from the totality of reading the comments.
5. The comments were then read again independently by two people, and scored as fitting a theme.
6. The two readers then compared their scoring of themes and any differences were resolved by discussion.
7. The themes were coded and then the additional information – chosen specialty, rejected specialty, whether the doctor was male or female, etc. – were added to the file of comments.
8. The records were analysed as those from doctors who specified that they had seriously considered, but then decided against trying for a career in (a) psychiatry and (b) all other specialties.

Appendix 2

Examples of statements about psychiatry

Job content

- 'Negative experience as a medical student'
- 'Unrewarding – difficult to make patients better'
- 'Not challenging enough – pace too slow'
- 'Fail to exercise majority of skills learnt as a medical student'
- 'Not enough medicine – waste of my learning'
- 'Realisation that practice different to theory'
- 'Psychiatry in small amounts is ok – but would probably be too much if full time'
- 'Experience in psychiatry as an F1 [Foundation Year 1, the first year after medical qualification] made me realise it was quite a narrow field that was too slow moving'
- 'Enjoyed the subject greatly, but had numerous bad placements that put me off'
- 'Violent patients'
- 'Singular dislike of personality disorders'

Self-appraisal

- 'Not suited to my personality'
- 'Too depressing'
- 'I feel physically scared of psychiatric patients'
- 'Too emotionally demanding'

References


- 1 Thomas K, Ellis A, Konrad T, Holzer C, Morrissey J. County-level estimates of mental health professional shortage in the United States. *Psychiatr Serv* 2009; **60**: 1323–28.

- 2 Tamaskar P, McGinnis RA. Declining student interest in psychiatry. *JAMA* 2002; **287**: 1859.
- 3 Salsberg E, Rockey PH, Rivers KL, Brotherton SE, Jackson GR. US residency training before and after the 1997 Balanced Budget Act. *JAMA* 2008; **300**: 1174–80.
- 4 Department of Health. NHS Vacancies Survey. Department of Health, 2003. <http://www.ic.nhs.uk/statistics-and-data-collections/workforce/nhs-and-gp-vacancies/nhs-vacancies-survey-england-31-march-2010>
- 5 Royal College of Psychiatrists. *Census 2009: Workforce Figures for Psychiatrists*. Royal College of Psychiatrists, 2010 (<http://www.rcpsych.ac.uk/PDF/2009%20Census.pdf>).
- 6 Goldacre MJ, Turner G, Fazel S, Lambert T. Career choices for psychiatry: national surveys of graduates of 1974–2000 from UK medical schools. *Br J Psychiatry* 2005; **186**: 158–64.
- 7 Lambert TW, Goldacre MJ, Edwards C, Parkhouse J. Career preferences of doctors who qualified in the United Kingdom in 1993 compared with those of doctors qualifying in 1974, 1977, 1980, and 1983. *BMJ* 1996; **313**: 19–24.
- 8 Goldacre MJ, Davidson JM, Lambert TW. Career choices at the end of the pre-registration year of doctors who qualified in the United Kingdom in 1996. *Med Educ* 1999; **33**: 882–9.
- 9 Goldacre M, Goldacre R, Lambert TW. Doctors who considered but did not pursue specific clinical specialties as careers: questionnaire surveys. *J R Soc Med* 2012; **105**: 166–76.
- 10 Sierles FS, Yager J, Weissman SH. Recruitment of U.S. medical graduates into psychiatry: reasons for optimism, sources of concern. *Acad Psychiatr* 2003; **27**: 252–9.
- 11 Taylor KS, Lambert TW, Goldacre MJ. Career progression and destinations, comparing men and women in the NHS: postal questionnaire surveys. *BMJ* 2009; **338**: b1735.
- 12 Goldacre MJ, Laxton L, Lambert TW. Medical graduates' early career choices of specialty and their eventual specialty destinations: UK prospective cohort studies. *BMJ* 2010; **340**: 1–9.
- 13 Eagles JM, Wilson S, Murdoch JM, Brown T. What impact do undergraduate experiences have upon recruitment into psychiatry? *Psychiatr Bull* 2007; **31**: 70–2.
- 14 Royal College of Psychiatrists. Revised action plan for the recruitment and retention of psychiatrists. Royal College of Psychiatrists, 2004 (available at <http://www.rcpsych.ac.uk/workforce.aspx>).
- 15 Collins J. *Foundation for Excellence: An Evaluation of the Foundation Programme*. Medical Education England, 2010.
- 16 Dein K, Livingston G, Bench C. 'Why did I become a psychiatrist?': survey of consultant psychiatrists. *Psychiatr Bull* 2007; **31**: 227–30.
- 17 Malhi GS, Parker GB, Parker K, Kirkby KC, Boyce P, Yellowlees P, et al. Shrinking away from psychiatry? A survey of Australian medical students' interest in psychiatry. *Aust NZ J Psychiatr* 2002; **36**: 416–23.
- 18 Lambert TW, Turner G, Fazel S, Goldacre M. Reasons why some UK medical graduates who initially choose psychiatry do not pursue it as a long-term career. *Psychol Med* 2006; **36**: 679–84.
- 19 Rajagopal S, Rehill KS, Godfrey E. Psychiatry as a career choice compared with other specialties: a survey of medical students. *Psychiatr Bull* 2004; **28**: 444–6.
- 20 Brown N, Vassilas CA, Oakley C. Recruiting psychiatrists – a Sisyphean task? *Psychiatr Bull* 2009; **33**: 390–2.
- 21 Katschnig H. Are psychiatrists an endangered species? Observations on internal and external challenges to the profession. *World Psychiatry* 2010; **9**: 21–8.
- 22 Centre for Workforce Intelligence. *Shape of the Medical Workforce: Informing Medical Training Numbers*. CFWI, 2011.
- 23 Fazel S, Ebmeier KP. Specialty choice in UK junior doctors: Is psychiatry the least popular specialty for UK and international medical graduates? *BMC Med Educ* 2009; **9**: 77.
- 24 Craddock N, Antebi D, Attenburrow M-J, Bailey A, Carson A, Cowen P, et al. Wake-up call for British psychiatry. *Br J Psychiatry* 2008; **193**: 6–9.
- 25 Sondergård L. Recruitment problems in psychiatry: just a matter of more exposition? *Acta Psychiatr Scand* 2007; **116**: 235–7.




Data supplement

(a) Men



(b) Women



(c) All




Fig. DS1 Percentages of responders from each cohort who chose psychiatry 1, 3, and 5 years after graduation (full colour bars show first choices, transparent bars above each full colour bar show second and third choices)

Table DS1 Numbers of respondents choosing psychiatry as their first, second or third choice of career at 1 (1974–2009), 3 (1974–2005) and 5 (1974–2000) years after graduation^a

Cohort	Numbers choosing psychiatry as first choice									Numbers choosing psychiatry as first, second or third choice								
	Year 1			Year 3			Year 5			Year 1			Year 3			Year 5		
	Men	Women	Total	Men	Women	Total	Men	Women	Total	Men	Women	Total	Men	Women	Total	Men	Women	Total
1974	54	26	80	38	21	59	48	29	77	123	62	185	58	34	92	60	44	104
1977	64	52	116	53	58	111	63	64	127	129	94	223	85	90	175	100	84	184
1980	65	40	105	94	56	150	85	64	149	137	83	220	131	86	217	112	83	195
1983	80	65	145	79	81	160	n/a ^b	n/a ^b	n/a ^b	168	158	326	123	130	253	n/a ^b	n/a ^b	n/a ^b
1993	59	57	116	53	71	124	58	79	137	96	103	199	75	106	181	66	96	162
1996	59	64	123	65	67	132	61	64	125	101	103	204	86	90	176	71	76	147
1999	53	76	129	50	80	130	55	89	144	78	115	193	58	116	174	65	96	161
2000	57	64	121	71	84	155	66	73	139	82	97	179	93	120	213	73	87	160
2002	50	66	116	38	76	114	41	65	106	77	103	180	47	98	145	42	70	112
2005	44	64	108	45	62	107	n/a ^b	n/a ^b	n/a ^b	73	111	184	63	82	145	n/a ^b	n/a ^b	n/a ^b
2008	56	86	142	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	88	142	230	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b
2009	33	89	122	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	57	160	217	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b	n/a ^b
Total	674	749	1423	586	656	1242	477	527	1004	1209	1331	2540	819	952	1771	589	636	1225

a. Numbers of respondents were: 16 488 (men), 17 486 (women) and 33 974 (total) in year 1; 13 429 (men), 12 745 (women) and 26 174 (total) in year 3; and 10 544 (men), 9 695 (women) and 20 239 (total) in year 5.

b. Third- and fifth-year choices not yet available for graduates of 2008/9; fifth-year choices not yet available for graduates of 2005; the 1983 cohort was not surveyed in year 5.

Table DS2 Comparison between doctors who chose psychiatry, general practice or other clinical specialties: numbers of doctors who specified that they were 'definite', 'probable' or 'uncertain' about their first choice of long-term career, at one, three, and five years after graduation^{a,b}

Firmness of choice	n								
	Year 1			Year 3			Year 5		
	Psychiatry	General practice	Other	Psychiatry	General practice	Other	Psychiatry	General practice	Other
Definite	423	3913	5152	669	5564	5884	649	5550	6997
Probable	677	4836	10 222	461	3220	6277	293	1778	2922
Uncertain	311	2096	4506	106	679	1836	49	288	483
Total	1411	10 845	19 880	1236	9463	13 997	991	7616	10 402

a. The numbers of respondents who did not answer this question were 12 (year 1), 8 (year 3) and 13 (year 5) of those choosing psychiatry; 62 (year 1), 33 (year 3) and 50 (year 5) of those choosing general practice; and 185 (year 1), 79 (year 3) and 72 (year 5) of those choosing other hospital specialties. Third- and fifth-year choices not yet available for graduates of 2008/9; fifth-year choices not yet available for graduates of 2005; the 1983 cohort was not surveyed in year 5.

b. Comparing psychiatry and general practice: Year 1 ($\chi^2_2=20.8$, $P<0.01$), Year 3 ($\chi^2_2=10.4$, $P<0.05$) and Year 5 ($\chi^2_2=23.8$, $P<0.001$). Comparing psychiatry and other hospital specialties: Year 1 ($\chi^2_2=11.6$, $P<0.01$), Year 3 ($\chi^2_2=71.6$, $P<0.001$), and Year 5 ($\chi^2_2=1.3$, $P=0.522$). Comparing men and women: Year 1 ($\chi^2_2=15.3$, $P<0.001$), Year 3 ($\chi^2_2=9.0$, $P<0.05$), and Year 5 ($\chi^2_2=1.9$, $P=0.393$).

Table DS3 Percentages and numbers of doctors from each medical school who specified psychiatry as a first, second or third choice of career at 1 (1993–2008), 3 (1993–2005) and 5 (1993–2002) years after graduation^a

Clinical medical school	%			n		
	Year 1	Year 3	Year 5	Year 1	Year 3	Year 5
England, older schools						
Birmingham	7.6	6.6	6.6	88	49	38
Bristol	5.1*	5.9	2.6**	42	34	12
Leeds	4.4**	4.5	4.4	41	31	25
Leicester	6.8	5.8	5.1	54	32	23
Liverpool	8.3	8.3*	7.7*	71	53	37
Manchester	6.0	5.6	5.1	91	60	44
Newcastle	6.5	5.8	5.0	63	39	27
Nottingham	4.9*	4.0*	4.5	47	26	24
Sheffield	6.2	6.3	6.0	62	43	31
Southampton	6.1*	6.6	6.1	48	38	28
Overall	6.2**	5.9	5.3	607	405	289
London						
Imperial College London	4.6**	4.9	4.6	61	49	39
King's College London	5.5	5.6	3.7*	84	63	34
Queen Mary, University of London & Westfield College	7.0	6.4	6.5	70	47	39
St George's	8.3	6.5	6.0	68	36	27
University College London	7.5	6.1	6.3	108	67	58
Overall	6.4*	5.8	5.3	391	262	197
Oxbridge						
Oxford	6.7	4.1	3.1	44	19	11
Cambridge	5.5	4.8	4.8	36	22	18
Overall	6.1	4.5*	4.0*	80	41	29
Scotland						
Aberdeen	10.0**	9.3	7.6*	74	54	34
Dundee	8.8*	7.1	3.8	57	33	14
Edinburgh	8.3*	8.6*	7.8**	96	73	54
Glasgow	7.1	8.6*	8.1**	76	71	55
Overall	8.4*	8.5**	7.2*	303	231	157
Northern Ireland						
Queens, Belfast	9.2*	8.5*	7.8*	76	52	39
Wales						
Cardiff/Wales unspecified	7.4	5.2	5.4	77	35	28
Total	6.8	6.3	5.6	1534	1026	739

** $P < 0.01$, * $P < 0.05$.

a. Logistic regression, excluding the new schools in England: after adjusting for gender and cohort differences, χ^2_{22} comparing the 23 remaining schools gave $P < 0.001$ for medical school differences in choices, in each of years 1, 3 and 5. Significant differences from the overall average in individual schools are denoted by ** for $P < 0.01$ and * for $P < 0.05$. An additional logistic regression model was used to investigate variation between medical school groupings ('overall' in the table), again excluding the new English schools. This model was also adjusted for gender and cohort. Years 1 and 3: χ^2_5 both $P < 0.001$; Year 5: $P < 0.01$. Year 3 and 5 choices not yet available for graduates of 2008/9; year 5 choices not yet available for graduates of 2005. Total numbers of respondents contributing to the table were 22 697 (1993–2009) in year 1, 16 289 (1993–2005) in year 3 and 13 109 (1993–2002) in year 5. Medical school was unknown for one respondent in year 1, for eight in year 3, and for three in year 5; these respondents are excluded. In the new schools in England (Brighton and Sussex, East Anglia, Hull York, Keele, Peninsula, and Warwick), 7.8% of students chose psychiatry in year 1; figures for individual schools are not quoted as they would be based on a single year of graduates.

Table DS4 Numbers of doctors who specified each factor as influencing their choice of long-term career a great deal: graduates of 1993, 1996, 1999, 2000, 2002, 2008 and 2009

Factor	Year 1			Year 3			Year 5		
	Psychiatry (n = 869)	General practice (n = 5615)	Other ^a (n = 12 829)	Psychiatry (n = 655)	General practice (n = 4020)	Other ^a (n = 8265)	Psychiatry (n = 651)	General practice (n = 4422)	Other ^a (n = 7448)
Domestic circumstances	213	2526	1852	81	1072	696	223	2789	1862
Hours/working conditions	473	3889	3433	198	1812	1553	396	3577	2735
Eventual financial prospects	66	657	1244	31	505	512	37	721	679
Promotion/career prospects	203	714	2500	96	503	1121	166	839	1747
Self-appraisal of own skills	450	2105	5159	228	1100	2280	440	2141	4055
Advice from others	75	682	1770	42	366	802	58	470	1074
Student experience of subject	474	1981	5778	125	461	1236	221	804	1701
Particular teacher/department ^b	164	585	3412	33	166	988	52	146	1274
Inclinations before medical school	158	668	1433	62	327	556	104	528	661
Experience of jobs so far	264	2082	5706	244	1197	3189	425	2166	5042
Enthusiasm/commitment: what I really want to do	612	3268	8713	230	1248	3062	486	2579	5609

a. Other hospital specialties (i.e. all hospital specialties other than psychiatry).
b. This statement was not presented to graduates of 1996 in their third year after graduating. Excluding this statement, the numbers of respondents to each statement varied between 699 and 808 (year 1), 336 and 343 (year 3) and 611 and 616 (year 5) of those choosing psychiatry; 4408 and 5179 (year 1), 2203 and 2223 (year 3) and 4063 and 4082 (year 5) of those choosing general practice; and 10421 and 12015 (year 1), 4553 and 4578 (year 3) and 6932 and 6980 (year 5) of those choosing other hospital specialties.

Table DS5 Number of doctors who gave each reason^a for rejecting psychiatry and for rejecting all other specialties combined, and total number of doctors (n)

Reason for rejection of specialty	Psychiatry (n = 99)	Other specialties (n = 2168)
Job content	71	722
Self-appraisal	12	90
Too stressful/lack of support/working conditions	8	224
Work-life balance	5	908
Training	2	181
Training too long	2	114
Future of the specialty uncertain/unstable	2	43
Advice	2	31
Working relationships	2	113
Lack of exposure and opportunities so far	2	45
Competition	1	275
Inadequate income	1	27
Personal circumstances	1	33
Fear of litigation	0	54

a. Some doctors gave more than one reason and we included each reason that applied.

BJPpsych

The British Journal of Psychiatry

Choice and rejection of psychiatry as a career: surveys of UK medical graduates from 1974 to 2009

Michael J. Goldacre, Seena Fazel, Fay Smith and Trevor Lambert

BJP published online October 25, 2012 Access the most recent version at DOI:

[10.1192/bjp.bp.112.111153](https://doi.org/10.1192/bjp.bp.112.111153)

Supplementary Material	Supplementary material can be found at: http://bjp.rcpsych.org/content/suppl/2012/10/24/bjp.bp.112.111153.DC1
References	This article cites 0 articles, 0 of which you can access for free at: http://bjp.rcpsych.org/content/early/2012/10/23/bjp.bp.112.111153#BIBL
Reprints/permissions	To obtain reprints or permission to reproduce material from this paper, please write to permissions@rcpsych.ac.uk
P<P	Published online 2012-10-25T00:05:33-07:00 in advance of the print journal.
You can respond to this article at	/letters/submit/bjprcpsych;bjp.bp.112.111153v1
Downloaded from	http://bjp.rcpsych.org/ on September 23, 2017 Published by The Royal College of Psychiatrists

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.